

The Abundance of Dwarf Galaxies

Hot gas explodes out of
young dwarf galaxies

Simulation by **Andrew Pontzen, Fabio Governato** and
Alyson Brooks on the **Darwin Supercomputer**, Cambridge UK.

Simulation code **Gasoline** by **James Wadsley** and **Tom Quinn**
with metal cooling by **Sijing Sheng**.

Visualization by **Andrew Pontzen**.

Alyson Brooks
Rutgers, the State University of New Jersey

In collaboration with the University of Washington's N-body Shop™
makers of quality galaxies

The Abundance of Dwarf Galaxies

Hot gas explodes out of
young dwarf galaxies

Simulation by **Andrew Pontzen, Fabio Governato** and
Alyson Brooks on the **Darwin Supercomputer**, Cambridge UK.

Simulation code **Gasoline** by **James Wadsley** and **Tom Quinn**
with metal cooling by **Sijing Sheng**.

Visualization by **Andrew Pontzen**.

Alyson Brooks

Rutgers, the State University of New Jersey

In collaboration with the University of Washington's N-body Shop™
makers of quality galaxies

The Abundance of Dwarf Galaxies

(1) BROOKS, PAPASTERGIS, ET AL., (2017), APJ,
SUBMITTED, ARXIV:1701.07835

**“HOW TO RECONCILE THE OBSERVED VELOCITY
FUNCTION OF GALAXIES WITH THEORY”**

(2) MUNSHI, BROOKS, ET AL., (2017), MNRAS,
SUBMITTED, ARXIV: TONIGHT!

**“GOING, GOING, GONE DARK: QUANTIFYING THE
SCATTER IN THE FAINTEST DWARF GALAXIES”**

Alyson Brooks

Rutgers, the State University of New Jersey

In collaboration with the University of Washington’s N-body Shop™
makers of quality galaxies

The Abundance of Dwarf Galaxies

**(1) BROOKS, PAPASTERGIS, ET AL., (2017), APJ,
SUBMITTED, ARXIV:1701.07835**

**“HOW TO RECONCILE THE OBSERVED VELOCITY
FUNCTION OF GALAXIES WITH THEORY”**

**(2) MUNSHI, BROOKS, ET AL., (2017), MNRAS,
SUBMITTED, ARXIV: TONIGHT!**

**“GOING, GOING, GONE DARK: QUANTIFYING THE
SCATTER IN THE FAINTEST DWARF GALAXIES”**


Alyson Brooks

Rutgers, the State University of New Jersey




In collaboration with the University of Washington’s N-body Shop™
makers of quality galaxies

THE MISSING DWARF PROBLEM IN THE FIELD

Observed number of galaxies






BUT: TWO WAYS TO MEASURE ROTATION (RESOLVED VS UNRESOLVED)




Observations


CREATING MOCK OBSERVATIONS




HOW WELL DO THEORY AND OBSERVATION MATCH?





NO MISSING DWARFS: WE OBSERVE THEM AT LOWER VELOCITIES THAN EXPECTED





NO MISSING DWARFS: WE OBSERVE THEM AT LOWER VELOCITIES THAN EXPECTED





WHY THE VELOCITY SHIFT?




WHY THE VELOCITY SHIFT?




WHY THE VELOCITY SHIFT?



WHY THE VELOCITY SHIFT?



POINT: ABUNDANCES ARE AS EXPECTED IN LCDM



The Abundance of Dwarf Galaxies


(1) BROOKS, PAPASTERGIS, ET AL., (2017), APJ,
SUBMITTED, ARXIV:1701.07835

**“HOW TO RECONCILE THE OBSERVED VELOCITY
FUNCTION OF GALAXIES WITH THEORY”**


(2) MUNSHI, BROOKS, ET AL., (2017), MNRAS,
SUBMITTED, ARXIV: TONIGHT!


**“GOING, GOING, GONE DARK: QUANTIFYING THE
SCATTER IN THE FAINTEST DWARF GALAXIES”**

THE STELLAR MASS — HALO MASS RELATION AT LOW MASSES




THE STELLAR MASS — HALO MASS RELATION







$z=0$ DM density



$z=0$ Gas density



$z=0$ DM density



$z=0$ Gas density

THE DWARF VOLUMES

Captain Marvel



COMPLETE

THE DWARF VOLUMES


Captain Marvel



Elektra





Storm




Rogue


COMPLETE







1-to-1 assumption of Abundance Matching break down at the low mass end!







THE “BEND” IS DUE TO THE INCLUSION OF SATELLITES







EFFECT OF SCATTER



A TOOL TO POPULATE LOW MASS HALOS




We adopt the trends defined for our well-resolved halos:

a slope of the SMHM $\alpha = 2.4$ for $8.4 < \log(M_{\text{halo}}) < 11.5$ (shown by the red dashed line), with a shallower slope of $\alpha = 0.64$ at lower halo masses. The scatter is fixed to a constant 0.2 above $M_1 = 5 \times 10^9 \text{ M}_{\odot}$, and linearly grows below this halo mass so that $\gamma = -0.26$.

$$\sigma = 0.2 + \gamma(\log_{10} M_{\text{halo}} - \log_{10} M_1)$$


PREDICTED STELLAR MASS FUNCTION




CONCLUSIONS

- Starting from the *abundance* of dwarf galaxies predicted in LCDM, the HIVF can be recovered. There is no missing dwarf problem in the field.
- The scatter in the SMHM relation in low mass halos increases with decreasing halo mass. There is no one-to-one assignment of stellar mass to halo mass.
- The “bend” at low halo masses found by Sawala et al. (2015) is due to the inclusion of satellites.
- The increased scatter at low masses leads to a prediction of a steeper stellar mass function in the ultra-faint dwarf galaxy mass range, currently being probed by DES, HSC, MagLiteS, etc, and by LSST in the future.

ALSO CONSIDER DETECTABILITY



NO MISSING DWARFS: WE OBSERVE THEM AT LOWER VELOCITIES THAN EXPECTED



NO MISSING DWARFS: WE OBSERVE THEM AT LOWER VELOCITIES THAN EXPECTED

