

Lattice Gauge Theory for $\mathcal{N} = 4$ Super Yang–Mills

David Schaich (Syracuse)



Lattice Gauge Theory for the LHC and Beyond
KITP Santa Barbara, 16 September 2015

[arXiv:1411.0166](https://arxiv.org/abs/1411.0166), [arXiv:1505.03135](https://arxiv.org/abs/1505.03135), [arXiv:1508.00884](https://arxiv.org/abs/1508.00884) & more to come
with Simon Catterall, Poul Damgaard, Tom DeGrand and Joel Giedt

Plan

- Motivations for lattice supersymmetry in general
- Lattice formulation of $\mathcal{N} = 4$ supersymmetric Yang–Mills
(new $\mathcal{O}(a)$ -improved action & [public code](#))
- Latest results for static potential and Konishi anomalous dim.
(confront with perturbation theory, AdS/CFT, bootstrap)
- Prospects and future directions
(sign problem; superQCD in two & three dimensions)

Motivation: Why lattice supersymmetry

Much interesting physics in 4D susy gauge theories:

dualities, holography, confinement, conformality, BSM, ...

Lattice promises non-perturbative insights from first principles

We can brainstorm many potential susy applications:

- Compute Wilson loops, spectrum, scaling dimensions, etc., going beyond perturbation theory, holography, bootstrap, ...
- New non-perturbative tests of conjectured dualities
- Predict low-energy constants from dynamical susy breaking
- Validate or refine AdS/CFT-based models for QCD phase diagram, condensed matter systems, ...

Motivation: Why lattice supersymmetry

Much interesting physics in 4D susy gauge theories:

dualities, holography, confinement, conformality, BSM, ...

Lattice promises non-perturbative insights from first principles

We can brainstorm many potential susy applications:

- Compute Wilson loops, spectrum, scaling dimensions, etc., going beyond perturbation theory, holography, bootstrap, ...
- New non-perturbative tests of conjectured dualities
- Predict low-energy constants from dynamical susy breaking
- Validate or refine AdS/CFT-based models for QCD phase diagram, condensed matter systems, ...

Many ideas probably infeasible; relatively few have been explored

Obstruction: Why not lattice supersymmetry

Recall supersymmetry extends Poincaré symmetry

by spinorial generators Q_{α}^I and $\bar{Q}_{\dot{\alpha}}^I$ with $I = 1, \dots, \mathcal{N}$

The super-Poincaré algebra includes $\{Q_{\alpha}^I, \bar{Q}_{\dot{\alpha}}^J\} = 2\delta^{IJ}\sigma_{\alpha\dot{\alpha}}^{\mu}P_{\mu}$

but infinitesimal translations don't exist in discrete space-time

Explicitly broken supersymmetry \implies relevant susy-violating operators

Typically many such operators,

especially with scalar fields from matter multiplets or from $\mathcal{N} > 1$

Fine-tuning couplings / counterterms to restore supersymmetry

is generally not practical in numerical lattice calculations

Solution: Exact susy on the lattice

Rapid progress in recent years

In certain systems (some subset of) the susy algebra
can be exactly preserved at non-zero lattice spacing

Equivalent constructions obtained from orbifolding / deconstruction
and from “topological” twisting — cf. [arXiv:0903.4881](https://arxiv.org/abs/0903.4881) for review

In four dimensions these constructions pick out a unique system:
maximally supersymmetric Yang–Mills ($\mathcal{N} = 4$ SYM)

$\mathcal{N} = 4$ SYM is a particularly interesting theory

- AdS/CFT correspondence
- Testing ground for reformulations of scattering amplitudes
- Arguably simplest non-trivial field theory in four dimensions

Basic features of $\mathcal{N} = 4$ SYM

$\mathcal{N} = 4$ SYM is a particularly interesting theory

—AdS/CFT correspondence

—Testing ground for reformulations of scattering amplitudes

—Arguably simplest non-trivial field theory in four dimensions

- $SU(N)$ gauge theory with four fermions Ψ^I and six scalars Φ^I , all massless and in adjoint rep.
- Action consists of kinetic, Yukawa and four-scalar terms with coefficients related by symmetries
- Supersymmetric: 16 supercharges Q_{α}^I and $\bar{Q}_{\dot{\alpha}}^I$ with $I = 1, \dots, 4$
Fields and Q 's transform under global $SU(4) \simeq SO(6)$ R symmetry
- Conformal: β function is zero for any 't Hooft coupling $\lambda = g^2 N$

What is special about $\mathcal{N} = 4$ SYM

An intuitive picture of topological twisting

$$\begin{pmatrix} Q_{\alpha}^1 & Q_{\alpha}^2 & Q_{\alpha}^3 & Q_{\alpha}^4 \\ \bar{Q}_{\dot{\alpha}}^1 & \bar{Q}_{\dot{\alpha}}^2 & \bar{Q}_{\dot{\alpha}}^3 & \bar{Q}_{\dot{\alpha}}^4 \end{pmatrix} = Q + Q_{\mu}\gamma_{\mu} + Q_{\mu\nu}\gamma_{\mu}\gamma_{\nu} + \bar{Q}_{\mu}\gamma_{\mu}\gamma_5 + \bar{Q}\gamma_5 \\ \rightarrow Q + Q_a\gamma_a + Q_{ab}\gamma_a\gamma_b \\ \text{with } a, b = 1, \dots, 5$$

Q 's transform with **integer spin** under “twisted rotation group”

$$\text{SO}(4)_{tw} \equiv \text{diag} \left[\text{SO}(4)_{\text{euc}} \otimes \text{SO}(4)_R \right] \quad \text{SO}(4)_R \subset \text{SO}(6)_R$$

This change of variables gives a susy subalgebra $\{Q, Q\} = 2Q^2 = 0$

This subalgebra can be exactly preserved on the lattice

Twisted $\mathcal{N} = 4$ SYM fields and \mathcal{Q}

Everything transforms with **integer spin** under $SO(4)_{tw}$ — **no spinors**

$$Q_{\alpha}^I \text{ and } \bar{Q}_{\dot{\alpha}}^I \longrightarrow \mathcal{Q}, \mathcal{Q}_a \text{ and } \mathcal{Q}_{ab}$$

$$\Psi^I \text{ and } \bar{\Psi}^I \longrightarrow \eta, \psi_a \text{ and } \chi_{ab}$$

$$A_{\mu} \text{ and } \Phi^{IJ} \longrightarrow \mathcal{A}_a = (A_{\mu}, \phi) + i(B_{\mu}, \bar{\phi}) \text{ and } \bar{\mathcal{A}}_a$$

Complexify gauge fields since scalars \longrightarrow vectors under twisting
(complexification $\implies U(N) = SU(N) \otimes U(1)$ gauge invariance)

The motivation is most obvious in five dimensions where

$$SO(5)_{tw} \equiv \text{diag} \left[SO(5)_{\text{euc}} \otimes SO(5)_R \right]$$

Then dimensional reduction takes gauge fields $A_a \longrightarrow (A_{\mu}, \phi)$
and scalar fields $B_a \longrightarrow (B_{\mu}, \bar{\phi})$

Twisted $\mathcal{N} = 4$ SYM fields and \mathcal{Q}

Everything transforms with **integer spin** under $SO(4)_{tw}$ — **no spinors**

$$Q_{\alpha}^I \text{ and } \bar{Q}_{\dot{\alpha}}^I \longrightarrow \mathcal{Q}, \mathcal{Q}_a \text{ and } \mathcal{Q}_{ab}$$

$$\Psi^I \text{ and } \bar{\Psi}^I \longrightarrow \eta, \psi_a \text{ and } \chi_{ab}$$

$$A_{\mu} \text{ and } \Phi^{IJ} \longrightarrow \mathcal{A}_a = (A_{\mu}, \phi) + i(B_{\mu}, \bar{\phi}) \text{ and } \bar{\mathcal{A}}_a$$

The twisted-scalar supersymmetry \mathcal{Q}

correctly interchanges bosonic \longleftrightarrow fermionic d.o.f. with $\mathcal{Q}^2 = 0$

$$\mathcal{Q} \mathcal{A}_a = \psi_a$$

$$\mathcal{Q} \psi_a = 0$$

$$\mathcal{Q} \chi_{ab} = -\bar{\mathcal{F}}_{ab}$$

$$\mathcal{Q} \bar{\mathcal{A}}_a = 0$$

$$\mathcal{Q} \eta = d$$

$$\mathcal{Q} d = 0$$

\nwarrow bosonic auxiliary field with e.o.m. $d = \bar{\mathcal{D}}_a \mathcal{A}_a$

Lattice $\mathcal{N} = 4$ SYM

The lattice theory is nearly a direct transcription,
despite breaking the 15 Q_a and Q_{ab}

- Covariant derivatives \longrightarrow finite difference operators
- Complexified gauge fields $\mathcal{A}_a \longrightarrow$ gauge links $\mathcal{U}_a \in \mathfrak{gl}(N, \mathbb{C})$

$$\begin{aligned} Q \mathcal{A}_a &\longrightarrow Q \mathcal{U}_a = \psi_a & Q \psi_a &= 0 \\ Q \chi_{ab} &= -\overline{\mathcal{F}}_{ab} & Q \overline{\mathcal{A}}_a &\longrightarrow Q \overline{\mathcal{U}}_a = 0 \\ Q \eta &= d & Q d &= 0 \end{aligned}$$

Geometry manifest: η and d on sites, \mathcal{U}_a and ψ_a on links, etc.


- Supersymmetric lattice action ($QS = 0$)
follows from $Q^2 \cdot = 0$ and **Bianchi identity**

$$S = \frac{N}{2\lambda_{\text{lat}}} Q \left(\chi_{ab} \mathcal{F}_{ab} + \eta \overline{\mathcal{D}}_a \mathcal{U}_a - \frac{1}{2} \eta d \right) - \frac{N}{8\lambda_{\text{lat}}} \epsilon_{abcde} \chi_{ab} \overline{\mathcal{D}}_c \chi_{de}$$

Five links in four dimensions $\longrightarrow A_4^*$ lattice


Revisit dimensional reduction in discrete spacetime,
treating all five \mathcal{U}_a symmetrically

- Start with hypercubic lattice
in 5d momentum space
- Symmetric** constraint $\sum_a \partial_a = 0$
projects to 4d momentum space
- Result is A_4 lattice
 \longrightarrow dual A_4^* lattice in real space



Twisted SO(4) symmetry on the A_4^* lattice

- Can picture A_4^* lattice as 4d analog of 2d triangular lattice
- Basis vectors are linearly dependent and non-orthogonal $\rightarrow \lambda = \lambda_{\text{lat}}/\sqrt{5}$
- Preserves S_5 point group symmetry



S_5 irreps precisely match onto irreps of twisted $\text{SO}(4)_{tw}$

$$\mathbf{5} = \mathbf{4} \oplus \mathbf{1} : \psi_a \rightarrow \psi_\mu, \bar{\eta}$$

$$\mathbf{10} = \mathbf{6} \oplus \mathbf{4} : \chi_{ab} \rightarrow \chi_{\mu\nu}, \bar{\psi}_\mu$$

$S_5 \rightarrow \text{SO}(4)_{tw}$ in continuum limit restores the rest of Q_a and Q_{ab}

Twisted $\mathcal{N} = 4$ SYM on the A_4^* lattice

High degree of exact symmetry: gauge invariance + \mathcal{Q} + \mathcal{S}_5

Several important analytic consequences:

- Moduli space preserved to all orders of lattice perturbation theory
→ no scalar potential induced by radiative corrections
- β function vanishes at one loop in lattice perturbation theory
- Real-space RG blocking transformations preserve \mathcal{Q} and \mathcal{S}_5
- Only one log. tuning to recover \mathcal{Q}_a and \mathcal{Q}_{ab} in the continuum

Not quite suitable for numerical calculations

Exact zero modes and flat directions must be regulated,
especially important in $U(1)$ sector

$\mathcal{N} = 4$ SYM lattice action (I)

$$S = \frac{N}{2\lambda_{\text{lat}}} Q \left(\chi_{ab} \mathcal{F}_{ab} + \eta \bar{D}_a \mathcal{U}_a - \frac{1}{2} \eta d \right) - \frac{N}{8\lambda_{\text{lat}}} \epsilon_{abcde} \chi_{ab} \bar{D}_c \chi_{de} + \mu^2 V$$


Scalar potential $V = \frac{1}{2N\lambda_{\text{lat}}} (\text{Tr} [\mathcal{U}_a \bar{\mathcal{U}}_a] - N)^2$ lifts SU(N) flat directions
and ensures $\mathcal{U}_a = \mathbb{I}_N + \mathcal{A}_a$ in continuum limit

Breaks Q **softly** — susy breaking automatically vanishes as $\mu^2 \rightarrow 0$

Violations of Ward identities $\langle Q\mathcal{O} \rangle = 0$
show Q breaking and restoration

Here considering

$$Q [\eta \mathcal{U}_a \bar{\mathcal{U}}_a] = d \mathcal{U}_a \bar{\mathcal{U}}_a - \eta \psi_a \bar{\mathcal{U}}_a$$



$$S = \frac{N}{2\lambda_{\text{lat}}} \mathcal{Q} \left(\chi_{ab} \mathcal{F}_{ab} + \downarrow - \frac{1}{2} \eta d \right) - \frac{N}{8\lambda_{\text{lat}}} \epsilon_{abcde} \chi_{ab} \bar{\mathcal{D}}_c \chi_{de} + \mu^2 V$$


$$\eta \left(\bar{\mathcal{D}}_a \mathcal{U}_a + G \sum_{a < b} [\det \mathcal{P}_{ab} - 1] \mathbb{I}_N \right)$$

Constraint on **plaquette det.** lifts U(1) zero mode & flat directions

\mathcal{Q} -exact implementation as new moduli space condition

Leads to $\langle \mathcal{Q}\mathcal{O} \rangle \propto (a/L)^2$,
much better than **naive constraint**

Looks like $\mathcal{O}(a)$ improvement
(\mathcal{Q} forbids all dim-5 operators)



Advertisement: Public code for lattice $\mathcal{N} = 4$ SYM

so that the full improved action becomes

$$\begin{aligned} S_{\text{imp}} &= S'_{\text{exact}} + S_{\text{closed}} + S'_{\text{soft}} \tag{3.10} \\ S'_{\text{exact}} &= \frac{N}{2\lambda_{\text{lat}}} \sum_n \text{Tr} \left[-\overline{\mathcal{F}}_{ab}(n) \mathcal{F}_{ab}(n) - \chi_{ab}(n) \mathcal{D}_{[a}^{(+)} \psi_{b]}(n) - \eta(n) \overline{\mathcal{D}}_a^{(-)} \psi_a(n) \right. \\ &\quad \left. + \frac{1}{2} \left(\overline{\mathcal{D}}_a^{(-)} \mathcal{U}_a(n) + G \sum_{a \neq b} (\det \mathcal{P}_{ab}(n) - 1) \mathbb{I}_N \right)^2 \right] - S_{\text{det}} \\ S_{\text{det}} &= \frac{N}{2\lambda_{\text{lat}}} G \sum_n \text{Tr} [\eta(n)] \sum_{a \neq b} [\det \mathcal{P}_{ab}(n)] \text{Tr} [\mathcal{U}_b^{-1}(n) \psi_b(n) + \mathcal{U}_a^{-1}(n + \widehat{\mu}_b) \psi_a(n + \widehat{\mu}_b)] \\ S_{\text{closed}} &= -\frac{N}{8\lambda_{\text{lat}}} \sum_n \text{Tr} [\epsilon_{abcde} \chi_{de}(n + \widehat{\mu}_a + \widehat{\mu}_b + \widehat{\mu}_c) \overline{\mathcal{D}}_c^{(-)} \chi_{ab}(n)], \\ S'_{\text{soft}} &= \frac{N}{2\lambda_{\text{lat}}} \mu^2 \sum_n \sum_a \left(\frac{1}{N} \text{Tr} [\mathcal{U}_a(n) \overline{\mathcal{U}}_a(n)] - 1 \right)^2 \end{aligned}$$

The full $\mathcal{N} = 4$ SYM lattice action is somewhat complicated
($\gtrsim 100$ gathers in the fermion operator)

To reduce barriers to entry our parallel code is publicly developed at
github.com/daschaich/susy

Evolved from MILC lattice QCD code, presented in [arXiv:1410.6971](https://arxiv.org/abs/1410.6971)

Application: Static potential is Coulombic at all λ

Static potential $V(r)$ from $r \times T$ Wilson loops: $W(r, T) \propto e^{-V(r) T}$


Fit $V(r)$ to Coulombic
or confining form

$$V(r) = A - C/r$$

$$V(r) = A - C/r + \sigma r$$

C is Coulomb coefficient

σ is string tension




Fits to confining form always produce vanishing string tension $\sigma = 0$

More sophisticated analyses in development using improved action

Coupling dependence of Coulomb coefficient

Continuum perturbation theory predicts $C(\lambda) = \lambda/(4\pi) + \mathcal{O}(\lambda^2)$

AdS/CFT predicts $C(\lambda) \propto \sqrt{\lambda}$ for $N \rightarrow \infty$, $\lambda \rightarrow \infty$, $\lambda \ll N$



$N = 2$ results agree with perturbation theory for all $\lambda \lesssim N$

$N = 3$ results bend down for $\lambda \gtrsim 1$ — approaching AdS/CFT?

Application: Konishi operator scaling dimension

$\mathcal{N} = 4$ SYM is conformal at all λ

→ power-law decay for all correlation functions

The Konishi operator is the simplest conformal primary operator

$$\mathcal{O}_K = \sum_I \text{Tr} [\Phi^I \Phi^I] \quad C_K(r) \equiv \mathcal{O}_K(x+r) \mathcal{O}_K(x) \propto r^{-2\Delta_K}$$

There are many predictions for its scaling dim. $\Delta_K(\lambda) = 2 + \gamma_K(\lambda)$

- From weak-coupling perturbation theory,
related to strong coupling by $\frac{4\pi N}{\lambda} \longleftrightarrow \frac{\lambda}{4\pi N}$ S duality
- From holography for $N \rightarrow \infty$ and $\lambda \rightarrow \infty$ but $\lambda \ll N$
- Upper bounds from the conformal bootstrap program

Only lattice gauge theory can access nonperturbative λ at moderate N

Konishi operator on the lattice

Extract scalar fields from polar decomposition of complexified links

$$U_a \simeq U_a (\mathbb{I}_N + \varphi_a)$$


$$\hat{O}_K = \sum_a \text{Tr} [\varphi_a \varphi_a]$$

$$\bar{O}_K = \hat{O}_K - \langle \hat{O}_K \rangle$$

$$\bar{C}_K(r) = \bar{O}_K(x+r) \bar{O}_K(x) \propto r^{-2\Delta_K}$$

Obvious sensitivity to volume
as desired for conformal system

Good tools to find Δ_K :
—Finite-size scaling
—Monte Carlo RG



Need lattice RG blocking scheme to carry out MCRG...

Real-space RG for lattice $\mathcal{N} = 4$ SYM

Lattice RG blocking transformation must preserve symmetries

Q and $S_5 \longleftrightarrow$ geometric structure of the system

Simple scheme constructed in [arXiv:1408.7067](https://arxiv.org/abs/1408.7067):

$$\begin{aligned} \mathcal{U}'_a(x') &= \xi \mathcal{U}_a(x) \mathcal{U}_a(x + \hat{\mu}_a) & \eta'(x') &= \eta(x) \\ \psi'_a(x') &= \xi [\psi_a(x) \mathcal{U}_a(x + \hat{\mu}_a) + \mathcal{U}_a(x) \psi_a(x + \hat{\mu}_a)] & & \text{etc.} \end{aligned}$$

Doubles lattice spacing $a \longrightarrow a' = 2a$, with ξ a tunable rescaling factor

Set ξ by equating plaquette on n -times-blocked L^4 ensemble
with that on independent $(n - 1)$ -times-blocked $(L/2)^4$ ensemble

Q -preserving RG blocking is necessary ingredient in derivation that
only one log. tuning needed to recover Q_a and Q_{ab} in the continuum

Scaling dimensions from MCRG stability matrix

Write system as (infinite) sum of operators, $H = \sum_i c_i \mathcal{O}_i$
with couplings c_i that flow under RG blocking transformation R_b

n -times-blocked system is $H^{(n)} = R_b H^{(n-1)} = \sum_i c_i^{(n)} \mathcal{O}_i^{(n)}$

Fixed point defined by $H^* = R_b H^*$ with couplings c_i^*

Linear expansion around fixed point defines **stability matrix** T_{ij}^*

$$c_i^{(n)} - c_i^* = \sum_j \left. \frac{\partial c_i^{(n)}}{\partial c_j^{(n-1)}} \right|_{H^*} (c_j^{(n-1)} - c_j^*) \equiv \sum_j T_{ij}^* (c_j^{(n-1)} - c_j^*)$$

Correlators of $\mathcal{O}_i, \mathcal{O}_j \longrightarrow$ elements of stability matrix (Swendsen, 1979)


Eigenvalues of $T_{ij}^* \longrightarrow$ scaling dimensions of corresponding operators

Preliminary Δ_K results from Monte Carlo RG

Far from bootstrap bounds

Aim to distinguish
perturbative vs. free theory

Rough agreement
between $N = 2, 3, 4$



Only statistical uncertainties so far, averaged over

★ 1 & 2 RG blocking steps

★ Blocked volumes 3^4 through 8^4

★ 1–5 operators in stability matrix

More sophisticated analyses in development,
while running larger volumes at stronger couplings

Practical question: Potential sign problem

In lattice gauge theory we compute operator expectation values

$$\langle \mathcal{O} \rangle = \frac{1}{\mathcal{Z}} \int [dU][d\bar{U}] \mathcal{O} e^{-S_B[U, \bar{U}]} \text{pf } \mathcal{D}[U, \bar{U}]$$

Pfaffian can be complex for lattice $\mathcal{N} = 4$ SYM, $\text{pf } \mathcal{D} = |\text{pf } \mathcal{D}| e^{i\alpha}$

Complicates interpretation of $\{e^{-S_B} \text{pf } \mathcal{D}\}$ as Boltzmann weight

We carry out phase-quenched RHMC, $\text{pf } \mathcal{D} \longrightarrow |\text{pf } \mathcal{D}|$

In principle need to reweight phase-quenched (pq) observables:



$$\langle \mathcal{O} \rangle = \frac{\langle \mathcal{O} e^{i\alpha} \rangle_{pq}}{\langle e^{i\alpha} \rangle_{pq}} \quad \text{with} \quad \langle \mathcal{O} e^{i\alpha} \rangle_{pq} = \frac{1}{\mathcal{Z}_{pq}} \int [dU][d\bar{U}] \mathcal{O} e^{i\alpha} e^{-S_B} |\text{pf } \mathcal{D}|$$

\implies Monitor $\langle e^{i\alpha} \rangle_{pq}$ as function of volume, coupling, N

Pfaffian phase dependence on volume and coupling

Left: $1 - \langle \cos(\alpha) \rangle_{pq} \ll 1$ independent of volume and N at $\lambda_{\text{lat}} = 1$

Right: New 4^4 results at $4 \leq \lambda_{\text{lat}} \leq 8$ show much larger fluctuations



Currently filling in more volumes and N for improved action

Extremely expensive analysis despite new parallel algorithm:

$\mathcal{O}(n^3)$ scaling $\longrightarrow \sim 50$ hours for single 4^4 measurement


Two puzzles posed by the sign problem

- With **periodic temporal boundary conditions** for the fermions we have an obvious sign problem, $\langle e^{i\alpha} \rangle_{pq}$ consistent with zero
- With **anti-periodic BCs** and all else the same $e^{i\alpha} \approx 1$, phase reweighting has negligible effect

Why such sensitivity to the BCs?

Also, other observables are nearly identical for these two ensembles

Why doesn't the sign problem affect other observables?



Recapitulation and outlook

Rapid recent progress in lattice supersymmetry

- Lattice promises non-perturbative insights from first principles
- Lattice $\mathcal{N} = 4$ SYM is practical thanks to exact \mathcal{Q} susy
- Public code to reduce barriers to entry

Latest results from ongoing calculations

- Static potential is Coulombic at all couplings, $C(\lambda)$ confronted with perturbation theory and AdS/CFT
- Promising initial Konishi anomalous dimension at weak coupling

Many more directions are being — or can be — pursued

- Understanding the (absence of a) sign problem
- Systems with less supersymmetry, in lower dimensions, including matter fields, exhibiting spontaneous susy breaking, ...

Thank you!

Thank you!

Collaborators

Simon Catterall, Poul Damgaard, Tom DeGrand and Joel Giedt

Funding and computing resources



Supplement: $(d - 1)$ -dimensional lattice superQCD

Method to add fundamental matter multiplets without breaking $Q^2 = 0$

Pioneered by Matsuura ([arXiv:0805.4491](https://arxiv.org/abs/0805.4491)), Sugino ([arXiv:0807.2683](https://arxiv.org/abs/0807.2683))

Consider 2-slice lattice

with $U(N) \times U(F)$ gauge group:


- (Adj, 1) fields on one slice
- (1, Adj) fields on the other
- Bi-fund. (\square , $\bar{\square}$) in between

Set $U(F)$ gauge coupling to zero

→ $U(N)$ in $d - 1$ dims.

with F fund. hypermultiplets

(Periodic BC → anti-fund.)





Spontaneous supersymmetry breaking

Can add Q -exact moduli space condition (Fayet–Iliopoulos D term),

$$\mathcal{S} \ni \eta \left(\overline{\mathcal{D}}_\mu \mathcal{U}_\mu + \sum_{i=1}^F \phi_i \overline{\phi}_i \right) \longrightarrow \eta \left(\overline{\mathcal{D}}_\mu \mathcal{U}_\mu + \sum_{i=1}^F \phi_i \overline{\phi}_i + r \mathbb{I}_N \right)$$

$\langle d \rangle = \left\langle \sum_{i=1}^F \phi_i \overline{\phi}_i + r \mathbb{I}_N \right\rangle \neq 0 \implies$ spontaneous susy breaking

Effectively $N \times N$ conditions imposed on $N \times F$ degrees of freedom...



Backup: Failure of Leibnitz rule in discrete space-time

Given that $\{Q_\alpha, \bar{Q}_{\dot{\alpha}}\} = 2\sigma_{\alpha\dot{\alpha}}^\mu P_\mu = 2i\sigma_{\alpha\dot{\alpha}}^\mu \partial_\mu$ is problematic,
why not try $\{Q_\alpha, \bar{Q}_{\dot{\alpha}}\} = 2i\sigma_{\alpha\dot{\alpha}}^\mu \nabla_\mu$ for a discrete translation?

Here $\nabla_\mu \phi(x) = \frac{1}{a} [\phi(x + a\hat{\mu}) - \phi(x)] = \partial_\mu \phi(x) + \frac{a}{2} \partial_\mu^2 \phi(x) + \mathcal{O}(a^2)$

Essential difference between ∂_μ and ∇_μ on the lattice, $a > 0$

$$\begin{aligned}\nabla_\mu [\phi(x)\chi(x)] &= a^{-1} [\phi(x + a\hat{\mu})\chi(x + a\hat{\mu}) - \phi(x)\chi(x)] \\ &= [\nabla_\mu \phi(x)] \chi(x) + \phi(x) \nabla_\mu \chi(x) + a [\nabla_\mu \phi(x)] \nabla_\mu \chi(x)\end{aligned}$$

We only recover the Leibnitz rule $\partial_\mu(fg) = (\partial_\mu f)g + f\partial_\mu g$ when $a \rightarrow 0$
 \implies “Discrete supersymmetry” breaks down on the lattice

(Dondi & Nicolai, “Lattice Supersymmetry”, 1977)

Backup: Twisting \longleftrightarrow Kähler–Dirac fermions

The Kähler–Dirac representation is related to the spinor $Q_\alpha^I, \bar{Q}_{\dot{\alpha}}^I$ by

$$\begin{pmatrix} Q_\alpha^1 & Q_\alpha^2 & Q_\alpha^3 & Q_\alpha^4 \\ \bar{Q}_{\dot{\alpha}}^1 & \bar{Q}_{\dot{\alpha}}^2 & \bar{Q}_{\dot{\alpha}}^3 & \bar{Q}_{\dot{\alpha}}^4 \end{pmatrix} = \mathcal{Q} + \mathcal{Q}_\mu \gamma_\mu + \mathcal{Q}_{\mu\nu} \gamma_\mu \gamma_\nu + \bar{\mathcal{Q}}_\mu \gamma_\mu \gamma_5 + \bar{\mathcal{Q}} \gamma_5 \\ \longrightarrow \mathcal{Q} + \mathcal{Q}_a \gamma_a + \mathcal{Q}_{ab} \gamma_a \gamma_b \\ \text{with } a, b = 1, \dots, 5$$


The 4×4 matrix involves R symmetry transformations along each row and (euclidean) Lorentz transformations along each column

\implies Kähler–Dirac components transform under “twisted rotation group”

$$\text{SO}(4)_{tw} \equiv \text{diag} \left[\text{SO}(4)_{\text{euc}} \otimes \text{SO}(4)_R \right] \\ \uparrow \\ \text{only } \text{SO}(4)_R \subset \text{SO}(6)_R$$

Backup: Hypercubic representation of A_4^* lattice

In the code it is very convenient to represent the A_4^* lattice as a hypercube with a backwards diagonal



Backup: More on flat directions

- 1 Complex gauge field $\implies U(N) = SU(N) \otimes U(1)$ gauge invariance
 $U(1)$ sector decouples only in continuum limit
- 2 $Q\mathcal{U}_a = \psi_a \implies$ gauge links must be elements of algebra
Resulting **flat directions** required by supersymmetric construction
but must be lifted to ensure $\mathcal{U}_a = \mathbb{I}_N + \mathcal{A}_a$ in continuum limit

We need to add two deformations to regulate flat directions

$$SU(N) \text{ scalar potential} \propto \mu^2 \sum_a (\text{Tr} [\mathcal{U}_a \bar{\mathcal{U}}_a] - N)^2$$

$$U(1) \text{ plaquette determinant} \sim G \sum_{a < b} (\det \mathcal{P}_{ab} - 1)$$

Scalar potential **softly** breaks Q supersymmetry

\swarrow susy-violating operators vanish as $\mu^2 \rightarrow 0$

Plaquette determinant can be made Q -invariant [[arXiv:1505.03135](https://arxiv.org/abs/1505.03135)]

Backup: One problem with flat directions



Gauge fields \mathcal{U}_a can move far away from continuum form $\mathbb{I}_N + \mathcal{A}_a$

if $\mu^2/\lambda_{\text{lat}}$ becomes too small

Example for $\mu = 0.2$ and $\lambda_{\text{lat}} = 5$ on $8^3 \times 24$ volume

Left: Bosonic action is stable $\sim 18\%$ off its supersymmetric value




Right: Polyakov loop wanders off to $\sim 10^9$



Backup: Another problem with U(1) flat directions

Can induce monopole condensation \longrightarrow transition to confined phase

This lattice phase is not present in continuum $\mathcal{N} = 4$ SYM



Around the same $\lambda_{\text{lat}} \approx 2 \dots$

Left: Polyakov loop falls towards zero

Center: Plaquette determinant falls towards zero

Right: Density of U(1) monopole world lines becomes non-zero

Backup: More on soft susy breaking



Before 2015 we used a more **naive constraint** on plaquette det.:

$$S_{soft} = \frac{N}{2\lambda_{lat}} \mu^2 \sum_a \left(\frac{1}{N} \text{Tr} [U_a \bar{U}_a] - 1 \right)^2 + \kappa \sum_{a < b} |\det P_{ab} - 1|^2$$

Both terms softly break \mathcal{Q} but $\det P_{ab}$ effects dominate

Left: The bosonic action provides another Ward identity $\langle s_B \rangle = 9N^2/2$

Right: Soft susy breaking is also suppressed $\propto 1/N^2$





Backup: More on supersymmetric constraints

arXiv:1505.03135 introduces method to impose \mathcal{Q} -invariant constraints

Basic idea: Modify aux. field equations of motion \rightarrow moduli space

$$d(n) = \overline{\mathcal{D}}_a^{(-)} \mathcal{U}_a(n) \rightarrow \overline{\mathcal{D}}_a^{(-)} \mathcal{U}_a(n) + \mathcal{GO}(n) \mathbb{I}_N$$

Putting both plaquette determinant and scalar potential in $\mathcal{O}(n)$
over-constrains system \rightarrow sub-optimal Ward identity violations





Backup: Code performance—weak and strong scaling

Results from [arXiv:1410.6971](https://arxiv.org/abs/1410.6971) for the pre-2015 (“unimproved”) action

Left: Strong scaling for U(2) and U(3) $16^3 \times 32$ RHMC

Right: Weak scaling for $\mathcal{O}(n^3)$ pfaffian calculation (fixed local volume)
 $n \equiv 16N^2L^3N_T$ is number of fermion degrees of freedom




Dashed lines are optimal scaling, solid line is power-law fit

Backup: Numerical costs for $N = 2, 3$ and 4 colors

Red: Find RHMC cost scaling $\sim N^5$ — recall adjoint fermion d.o.f. $\propto N^2$

Blue: Pfaffian cost scaling consistent with expected N^6

Additional factor of $\sim 2\times$ from new improved action




Backup: Restoration of Q_a and Q_{ab} supersymmetries

Results from [arXiv:1411.0166](https://arxiv.org/abs/1411.0166) to be revisited with the new action...

Q_a and Q_{ab} from restoration of R symmetry (motivation for A_4^* lattice)

Modified Wilson loops test R symmetries at non-zero lattice spacing

Parameter c_2 may need log. tuning in continuum limit




Backup: $\mathcal{N} = 4$ SYM static potential from Wilson loops

Extract static potential $V(r)$ from $r \times T$ Wilson loops

$$W(r, T) \propto e^{-V(r) T}$$

$$V(r) = A - C/r + \sigma r$$


Coulomb gauge trick from lattice QCD reduces A_4^* lattice complications



Backup: Perturbation theory for Coulomb coefficient

For range of couplings currently being studied

(continuum) perturbation theory for $C(\lambda)$ is well behaved





Backup: More tests of the U(2) static potential

Left: Projecting Wilson loops from U(2) \longrightarrow SU(2)

$$\implies \text{factor of } \frac{N^2-1}{N^2} = 3/4$$

Right: Unitarizing links removes scalars \implies factor of 1/2





Some results slightly above expected factors,
may be related to non-zero auxiliary couplings μ and κ / G

Backup: More tests of the U(3) static potential

Left: Projecting Wilson loops from U(3) \longrightarrow SU(3)

$$\implies \text{factor of } \frac{N^2-1}{N^2} = 8/9$$

Right: Unitarizing links removes scalars \implies factor of 1/2



Some results slightly above expected factors,
may be related to non-zero auxiliary couplings μ and κ / G

Backup: Smearing for Konishi analyses

As in glueball analyses, smearing enlarges operator basis

Using APE-like smearing: $(1 - \alpha) \square + \frac{\alpha}{8} \sum \square$,

with staples built from unitary parts of links but no final unitarization
(unitarized smearing — e.g. stout — doesn't affect Konishi)

Average plaquette is stable upon smearing (**right**)

while minimum plaquette steadily increases (**left**)

