

# ON SOLUTIONS OF MULTI-PARTON 'T HOOFT EQUATIONS

Jacek Wosiek  
Jagellonian University & KITP

In collaboration with Daniele Dorigoni and Gabriele Veneziano



INNOVATIVE ECONOMY  
NATIONAL COHESION STRATEGY



*Foundation for Polish Science*

EUROPEAN REGIONAL  
DEVELOPMENT FUND



# 1 Outline

- An alternative to lattice - diagonalize the Hamiltonian
- On the Light Front - numerics: Light Cone Discretization
- Simplifications (I):
  - large N - planar diagrams - single traces
  - less dimensions - reductions
  - even quantum mechanics (but at  $N \rightarrow \infty$ )
  - supersymmetry
- QCD equations: eigenequations for  $H_{LC}$ 
  - coupled Bethe-Salpeter equations on the LC
  - simplifications (II) - **Coulomb Approximation**
- 't Hooft equations with many partons
- Solutions – numerical
- Solutions – **analytical**

## 2 Diagonalizing Hamiltonian

### 2.1 One way: Light Cone Discretization

$$P^+ = \sum_{i=1}^n p_i^+, \quad p_i^+ > 0$$
$$K = \sum_{i=1}^n k_i, \quad K, k_i - \text{integer } (> 0),$$

Cutoff  $K \implies$  partitions  $\{k_1, k_2, \dots\} \implies$  states

$$|\{k\}\rangle = Tr[a^\dagger(k_1)a^\dagger(k_2)\dots a^\dagger(k_n)]|0\rangle \quad (1)$$

$$|\{k\}\rangle \implies \langle\{k\}|H|\{k'\}\rangle \implies E_n$$



## 2.2 Second way: integral equations in the continuum

- Different cutoff (on parton multiplicity) – directly in the continuum

$$H|\Phi\rangle = M^2|\Phi\rangle \tag{2}$$

$$|\Phi\rangle \rightarrow \Phi_n(x_1, x_2, \dots, x_n)$$

$\leftrightarrow$



$$M^2\Phi_n(x_1 \dots x_n) = A \otimes \Phi_n + B \otimes \Phi_{n-2} + C \otimes \Phi_{n+2} \tag{3}$$

• EQUATIONS

$$|\Phi\rangle = \sum_{n=2}^{\infty} \int [dx] \delta(1 - x_1 - x_2 - \dots - x_n) \Phi_n(x_1, x_2, \dots, x_n) \text{Tr}[a^\dagger(x_1) a^\dagger(x_2) \dots a^\dagger(x_n)] |0\rangle$$

EXAMPLE 1:  $QCD_2$  ( fundamental fermions )

$$M^2 f(x) = m^2 \left( \frac{1}{x} + \frac{1}{1-x} \right) f(x) + \frac{\lambda}{\pi} \int_0^1 dy \frac{f(x) - f(y)}{(y-x)^2}$$

$$f(x) = \Phi_2(x, 1-x)$$

EXAMPLE 2:  $SYM_2$  restricted to the two-parton sector

There are two coupled equations in the bosonic sector

$$M^2\phi_{bb}(x) = m_b^2 \left( \frac{1}{x} + \frac{1}{1-x} \right) \phi_{bb}(x) + \frac{\lambda}{2} \frac{\phi_{bb}(x)}{\sqrt{x(1-x)}} - \frac{2\lambda}{\pi} \int_0^1 \frac{(x+y)(2-x-y)}{4\sqrt{x(1-x)y(1-y)}} \frac{[\phi_{bb}(y) - \phi_{bb}(x)]}{(y-x)^2} dy + \frac{\lambda}{2\pi} \int_0^1 \frac{1}{(y-x)} \frac{\phi_{ff}(y)}{\sqrt{x(1-x)}} dy$$

$$M^2\phi_{ff}(x) = m_f^2 \left( \frac{1}{x} + \frac{1}{1-x} \right) \phi_{ff}(x) - \frac{2\lambda}{\pi} \int_0^1 \frac{[\phi_{ff}(y) - \phi_{ff}(x)]}{(y-x)^2} dy + \frac{\lambda}{2\pi} \int_0^1 \frac{1}{(x-y)} \frac{\phi_{bb}(y)}{\sqrt{y(1-y)}} dy$$

and the single one in the fermionic sector

$$M^2\phi_{bf}(x) = \left( \frac{m_b^2}{x} + \frac{m_f^2}{1-x} \right) \phi_{bf}(x) + \frac{2\lambda}{\pi} \frac{\phi_{bf}(x)}{\sqrt{x+x}} - \frac{2\lambda}{\pi} \int_0^1 \frac{(x+y)}{2\sqrt{xy}} \frac{[\phi_{bf}(y) - \phi_{bf}(x)]}{(y-x)^2} dy - \frac{\lambda}{2\pi} \int_0^1 \frac{1}{(1-y-x)} \frac{\phi_{bf}(y)}{\sqrt{xy}} dy$$

(4)

Example 3:  $YM_2$  with adjoint fermionic matter - all parton-number sectors

$$\begin{aligned}
M^2 \phi_n(x_1 \dots x_n) &= \frac{m^2}{x_1} \phi_n(x_1 \dots x_n) \\
&+ \frac{\lambda}{\pi} \frac{1}{(x_1 + x_2)^2} \int_0^{x_1+x_2} dy \phi_n(y, x_1 + x_2 - y, x_3 \dots x_n) \\
&+ \frac{\lambda}{\pi} \int_0^{x_1+x_2} \frac{dy}{(x_1 - y)^2} \{ \phi_n(x_1, x_2, x_3 \dots x_n) \\
&\quad - \phi_n(y, x_1 + x_2 - y, x_3 \dots x_n) \} \\
&+ \frac{\lambda}{\pi} \int_0^{x_1} dy \int_0^{x_1-y} dz \phi_{n+2}(y, z, x_1 - y - z, x_2 \dots x_n) \left[ \frac{1}{(y+z)^2} - \frac{1}{(x_1-y)^2} \right] \\
&+ \frac{\lambda}{\pi} \phi_{n-2}(x_1 + x_2 + x_3, x_4 \dots x_n) \left[ \frac{1}{(x_1 + x_2)^2} - \frac{1}{(x_1 - x_3)^2} \right] \\
&\pm \text{cyclic permutations of } (x_1 \dots x_n)
\end{aligned}$$

### 3 This work (JHEP 1106:051, 2011)

- $\mathcal{N} = 1$ ,  $SYM_4$  on the LC
- Reduce  $D = 4 \longrightarrow D = 2 \implies QCD_2$  with adjoined matter
- **The Coulomb Approximation - keep only most singular (IR) terms in H**
  1. diagonal in parton multiplicity – can study each  $p$  separately, here  $p = 2, 3, 4$
  2. eigenvalues – spectrum
  3. eigenstates – wave functions **also in  $x$  - space**
  4. confinement – determine string tension

## 4 Coulomb divergences

- IR divergences (logarithmic) couple different multiplicity sectors
- Coulomb divergences (linear), but they cancel within one multiplicity
- Can be done independently for each parton multiplicity  $p$

### A possibility

- $\longrightarrow$  Solve Coulomb problem first, and then successively add radiation

Simplified Hamiltonian  $SYM_4 \implies SYM_2 \implies H_{Coulomb}$

$$H_{Coulomb}^{quad} = \frac{\lambda}{\pi} \int_0^\infty dk \int_0^k \frac{dq}{q^2} \text{Tr}[A_k^\dagger A_k] \quad (5)$$

$$H_{Coulomb}^{quartic} = -\frac{g^2}{2\pi} \int_0^\infty dp_1 dp_2 \left[ \int_0^{p_1} \frac{dq}{q^2} \text{Tr}[A_{p_1}^\dagger B_{p_2}^\dagger B_{p_2+q} A_{p_1-q}] \right. \\ \left. + \int_0^{p_2} \frac{dq}{q^2} \text{Tr}(A_{p_2}^\dagger B_{p_1}^\dagger B_{p_1+q} A_{p_2-q}) \right]$$

## 5 Two partons

$$|k, K - k\rangle, \quad k = 1, \dots, K - 1 \quad (6)$$

$$\langle k|H|k'\rangle \Rightarrow |\Phi_n\rangle \Rightarrow \Phi_n(k) \xrightarrow{FT} \Phi_n(d_{12}) \quad (7)$$




Figure 1:  $\rho_n(d_{12})$ ,  $p = 2$ ,  $K = 200$ ,  $n = 1, 25, 50, 100, 150, 199$ .

# Linear spectrum for two partons




Figure 2: **Eigenenergies of the,  $p=2$ , excited states as a function of the relative separation between two partons,  $K = 30, 50, 100, 200$ .**

## 6 Three partons - generalization of the 't Hooft solution to many bodies

$$|k_1, k_2, K - k_1 - k_2\rangle, \quad k_1 = 1, \dots, K - 2, \quad k_2 = 1, \dots, K - k_1 - 1$$

$$\langle k_1, k_2 | H | k'_1, k'_2 \rangle \Rightarrow |\Phi_n\rangle \Rightarrow \Phi_n(k_1, k_2) \xrightarrow{FT} \Phi_n(d_{13}, d_{23})$$




Figure 3:  $\rho_1(d_{13}, d_{23})$




Figure 4:  $|\rho_{10}(d_{13}, d_{23})|$




Figure 5:  $\rho_{50}(d_{13}, d_{23})$




Figure 6:  $\rho_{100}(d_{13}, d_{23})$




Figure 7:  $\rho_{200}(d_{13}, d_{23})$




Figure 8:  $\rho_{300}(d_{13}, d_{23})$




Figure 9:  $\rho_{400}(d_{13}, d_{23})$

The highest state




Figure 10:  $\rho_{406}(d_{13}, d_{23})$

## And on the Dalitz plot




Figure 11: Series B. As above but on the Dalitz plot. Now diquarks are allowed,  $d_{min} = 0$

## Linear spectrum for three partons




Figure 12: Eigenenergies of the,  $p=3$ , excited states as a function of the combined length of strings stretching between three partons.

## Four partons




Figure 13: Structure of eigenstates with four partons. Contour plots in three relative distances  $(d_{14}, d_{24}, d_{34})$  for states no. 1,9,35,60,100,165 spanning the whole range of states for  $K = 12$ ,  $r_{max} = 165$ .

## 7 Analytic solutions

- Massless quarks

$$\frac{\lambda}{\pi} \int_0^P dk \frac{f(p) - f(k)}{(p - k)^2} = E_C f(p) \longrightarrow \text{Fig.2}$$

- Assume that the singularity dominates (e.g. for large  $E_C$ ) [Kutasov, '95]

$$\begin{aligned} \frac{\lambda}{\pi} \int_{-\infty}^{\infty} dk \frac{f(p) - f(k)}{(p - k)^2} &= E_C f(p) \\ f(k) = \exp(ik\Delta) &\longrightarrow E_C = \lambda|\Delta|, \quad \Delta = r_2 - r_1 \end{aligned} \quad (8)$$

- a generic solution -  $\Delta$  arbitrary

- boundary conditions

- massless quarks  $\longrightarrow$  Neumann:  $f'(0) = f'(P) = 0$  [Neuberger, '04]

$$\Delta = \frac{n2\pi}{2P} = \frac{n}{2}a$$

$$f_n(k) = \cos(\pi nk/P) = \cos(\pi nx_F) \quad [\text{'t Hooft, '74}]$$

## Two partons: numerics vs. analytics




Figure 14: Comparison of numerical (DLCQ) and analytical (WKB) results for the two LC wave functions in the two parton sector

## 8 Analytic solution in many parton sectors

- Strategy:

- general solution of the asymptotic equation for  $n$  partons

- derive boundary conditions (BC) for  $n$  partons

- identification of independent (and complete) set of solutions satisfying BC

- classifying solutions w.r.t. their behaviour under  $Z_n$

- n-parton 't Hooft equation

$$\frac{\lambda}{2\pi} \int_0^{p_1+p_2} dk \frac{\psi_n(p_1, p_2, p_3 \dots p_n) - \psi_n(k, p_1 + p_2 - k, p_3 \dots p_n)}{(p_1 - k)^2} \pm \text{cyclic permutations of } (p_1 \dots p_n) = E_C \psi_n(p_1 \dots p_n) \quad (9)$$

- phase space

$$p_1 + p_2 + \dots + p_n = P, \quad p_i > 0 \quad (10)$$

only  $n - 1$  independent momenta,

e.g. for  $n = 2$   $\psi_2(p_1, P - p_1) = f(p_1)$

- phase space boundaries:  $p_i = 0, \quad i = 1, \dots, n.$

- Boundary conditions - two partons

$$M^2 f(x) = m^2 \left( \frac{1}{x} + \frac{1}{1-x} \right) f(x) + \frac{\lambda}{\pi} PV \int_0^1 dy \frac{f(x) - f(y)}{(y-x)^2}$$

- $m > 0$   $\longrightarrow$  Dirichlet
- $m = 0$   $\longrightarrow$  Neumann
- BC for n massless partons: **generalization** of Neumann conditions

$$\begin{aligned} p_1 = 0 & : (\partial_2 - 2\partial_1)\psi = 0 \\ p_i = 0 & : (\partial_{i+1} - 2\partial_i + \partial_{i-1})\psi = 0, \quad 2 \leq i \leq n-2 \\ p_{n-1} = 0 & : (\partial_{n-2} - 2\partial_{n-1})\psi = 0 \\ p_n = 0 & : (\partial_1 + \partial_{n-1})\psi = 0 \end{aligned}$$

[ Z. Ambrozinski ]

BC follow from a requirement of cancellation of IR divergences at the boundaries of the phase space.

- generic solution of asymptotic ( $f_0^{x_i+x_j} \dots \longrightarrow f_{-\infty}^{\infty} \dots$ ) equations in n parton sector

$$\psi(k_1, \dots, k_n) = \exp(ik_1 r_1 + ik_2 r_2 + \dots + ik_n r_n) \quad (11)$$

- asymptotic eigenvalue

$$E_C = \frac{\lambda}{2} \sum_{i=1}^n |\Delta_{i,i+1}|, \quad \Delta_{i,j} = r_i - r_j, \quad n+1 = 1. \quad (12)$$

- How to construct solutions which satisfy BC ??

## 9 Three partons

- New feature of  $n > 2$  sectors: degeneracy  $\longrightarrow$  use more trial functions with the same eigenvalue

Sufficient set for  $n = 3$

$$\begin{aligned}
\Psi_1 &= \exp(+i(k_1 r_1 + k_2 r_2 + k_3 r_3)) \\
\Psi_2 &= \exp(-i(k_1 r_1 + k_3 r_2 + k_2 r_3)) \exp(i2Pr_1) \\
\Psi_3 &= \exp(+i(k_2 r_1 + k_3 r_2 + k_1 r_3)) \\
\Psi_4 &= \exp(-i(k_3 r_1 + k_2 r_2 + k_1 r_3)) \exp(i2Pr_2) \\
\Psi_5 &= \exp(+i(k_3 r_1 + k_1 r_2 + k_2 r_3)) \\
\Psi_6 &= \exp(-i(k_2 r_1 + k_1 r_2 + k_3 r_3)) \exp(i2Pr_3)
\end{aligned}$$

Or in terms of independent momenta and coordinate differences

$$\begin{aligned}
\psi_1 &= \exp(i(k_1 \Delta_{13} + k_2 \Delta_{23})) \exp(iPr_3) \\
\psi_2 &= \exp(i(k_1 \Delta_{21} + k_2 \Delta_{23})) \exp(iP(r_3 + \Delta_{13} + \Delta_{12})) \\
\psi_3 &= \exp(i(k_1 \Delta_{32} + k_2 \Delta_{12})) \exp(iP(r_3 + \Delta_{23})) \\
\psi_4 &= \exp(i(k_1 \Delta_{13} + k_2 \Delta_{12})) \exp(iP(r_3 + \Delta_{23} + \Delta_{21})) \\
\psi_5 &= \exp(i(k_1 \Delta_{21} + k_2 \Delta_{31})) \exp(iP(r_3 + \Delta_{13})) \\
\psi_6 &= \exp(i(k_1 \Delta_{32} + k_2 \Delta_{31})) \exp(iPr_3)
\end{aligned}$$

- Necessary condition for BC: on each plane some subsets have to have the same dependence on all other (not fixed) variables.

E.g. on  $k_1 = 0$  boundary cancellations may occur only within (1,2) , (3,4) and (5,6) pairs.

- **Indeed**, for integer (in units of  $2\pi/P$ )  $\Delta$ 's, all BC's are satisfied by

$$\psi_{r,s}(k_1, k_2) = \sum_{i=1}^6 \psi_i = \psi^{singlet}, \quad \Delta_{13} = \frac{r}{2}, \quad \Delta_{23} = \frac{s}{2}, \quad r, s \text{ even}$$

- $Z_3$  covariant solutions can be constructed as well

$$\psi_{r,s,\nu}(k_1, k_2) = \psi_1 + \lambda\psi_5 + \lambda^2\psi_3 + \psi_2 + \psi_4 + \psi_6$$

$$\Delta_{13} = \frac{r + \nu}{2}; \quad \Delta_{2,3} = \frac{s - \nu}{2} \quad \nu = \pm\frac{1}{3}, \quad \lambda = e^{2\pi i\nu}, \quad r, s \text{ odd.}$$

this quantization follows from

$$\exp(iP\Delta_{13}) = \lambda^2, \quad \exp(iP\Delta_{23}) = \lambda,$$

which generalizes the  $\exp(iP\Delta_{12}) = \pm 1$  from the two parton case.

- all pairs  $(r, s)$  generate overcomplete sets
- for a complete basis **it suffices** to use

$(r, s) = (2n, 2l)$  and/or  $(2l, 2n)$ ,  $0 \leq l \leq [n/2]$ .  
 for each eigenvalue  $E_C = \frac{\lambda}{2}La$  and  $\nu = 0$ ,  
 where the "combined length of strings"  $L = 2n$ .  
 $\longrightarrow$  each  $E_C(n)$  has degeneracy

$$g_n = \begin{cases} n + 1, & n \text{ even} \\ n, & n \text{ odd} \end{cases} \quad (13)$$

and for  $\nu = 1/3$  :

$$L^I = 2n + 1 + \nu, \quad L^{II} = 2n + 3 - \nu, \quad (14)$$

$$(r, s)^I = (2n + 1, 2l + 1), \quad (r, s)^{II} = (2l + 1, 2n + 3) \quad (15)$$

## 9.1 Comparison with numerical results

- Profiles of non degenerate states agree very well, c.f. Table 1 for  $\nu = 1/3$
- Eigenenergies differ by 50% for the lowest state.

The discrepancy goes down to 30% around  $no = 13 \leftrightarrow$  WKB.

<i>num. - no's</i>	<i>anal. - (r, s)</i>	$ \langle num   anal \rangle ^2$	$LP/2\pi$	$E_{anal}$	$E_{num}$
1	(0,0)	1.0	0	0	0
4	(2,2)	.96	2	39.5	22.0
(2,3)	(1,1)	.96	4/3	26.3	11.3
(5,6)	(1,3)	.93	8/3	52.6	29.3
(7,8)	(3,3)	.91	10/3	65.8	39.0
(12,13)	(3,5)	.87	14/3	92.1	58.2

Table 1: First six states in the  $\nu = 0, 1/3$  sector, comparison with numerical (DLCQ) calculations.

- for higher states (i.e. with degeneracy): analytical solutions with degeneracy  $g$  correspond uniquely to a group of  $g$  numerical eigenstates (substantial overlaps)




Figure 15: Correspondence between the numerical (left) and analytical (right) spectra. Only  $Z_3$  singlets are shown. Analytic levels are  $g$ -fold degenerate, here  $g=1,3,3,5,5$  and  $7$  respectively.  $\rho = 1.3$

- High eigenvalues - can test completeness and WKB by comparing the entropy, or rather the number of states with energy below  $E$ .




Figure 16: Energy distribuant  $N(E, 1/K)$  and its extrapolation to  $K = \infty$




Figure 17: Effective scale factor obtained from  $N_{num}(E, K = \infty) = N_{anal}(E/\rho)$

## 10 Four partons

- Trial states are direct generalization of symmetric sums from the  $n = 3$  case.
- They are characterized by a triple of integers  $(d_{12}, d_{23}, d_{34})$ ,  $d = \Delta P/2\pi$ .
- They **DO NOT** satisfy our boundary conditions !
- However their simple combinations DO .

## Procedure

1. Generate all sets of above triples which satisfy

$$\sum_i^4 |d_{i,i+1}| = L = 2n, \quad (16)$$

for a given  $n$ .

2. Identify linearly independent subset of corresponding trials
3. Search for the **linearly dependent combinations on the boundary planes** by inspecting generalized Wronskians of corresponding partial derivatives.
4. Identify combinations satisfying our boundary conditions.
5. Organize states found in pt. 4 by choosing some labeling scheme.
6. Check completeness of this basis as in the three parton case.

## Results

- A. Indeed a series of simple linear combinations, which satisfy boundary conditions (BC) on all boundary planes, exists.
- B. Only combinations, which appear, contain one (singles), two (doubles) and three (triples) basis functions from step (2).
- C. Each independent trial function from step (2) appears once and only once in one of the combinations. All independent trials are used.
- D. Relative coefficients of all combinations found are very simple: all 1's in triples, and 1 and 2 in doubles. This finds a nice explanation upon the detailed inspection below.
- E. All combinations are orthogonal even though the original basis, found in 2, was not.




Figure 18: Solutions with 4 partons on the  $(d_{12}, d_{23}) = (i, j)$  plane, together with the contour plots (blue) of  $|d_{12}| + |d_{23}| + |a - d_{12} - d_{23}| = 2n - |a|$  for fixed  $a = d_{12} + d_{23} + d_{34} = n, n - 1, n - 2, \dots; n=13$ . Reflections across the black lines provide triples which satisfy BC.





Figure 19:  $x$  profile: numeric (left) and analytic (right),  $y = z = 1.3$




Figure 20:  $(y, z)$  contour plots of the same profile: numeric vs. analytic as above,  $x = 0$




Figure 21: Scale factor for four partons

## 11 Arbitrary number of partons $p$

$p = 5$  - similar to  $p=4$ : trials, basis of independent solutions,

Wronskians  $\Rightarrow$  combinations which satisfy BC (more than triples: 4-,6-,12- plets)

$\Rightarrow$  Rules (emerged from analyzing  $p=4,5$ )

**Rule I** (to generate basis of trial solutions)

- generate all closed loops (made of  $p$  "bits") with size  $d$  and energy  $L$
- mod out  $Z_p$  and  $IZ_p$
- sum over  $d$  at fixed  $L$

**Rule II** (to construct combinations satisfying BC)

- Solutions with the same values of  $\{d's\}$  form combinations which satisfy BC's.  
e.g.  $(1, 0, 2, -3)$  and  $(0, 1, 2, -3)$  for  $p = 4$

## Counting states ( for $p \leq 6$ )




Figure 22: Entropy of solutions (vs.  $M^2/\lambda$ ) from the first six multiplicity sectors.

$$\rho(M) \sim \exp M/T_H, \quad T_H = \frac{1.6 - 1.7}{\sqrt{\pi}} \sqrt{\lambda} \leftrightarrow (1.3 - 1.4) \text{ [Bhanot, et.al ]}$$

## 12 Summary

- Need a string-like counting of states for arbitrary  $p > 4$
- Interpretation of  $T_H$  - confirmation with higher  $p$  ?
- Green's functions  $\longrightarrow$  solve the hierarchy by Gauss elimination !
- Add transverse degrees of freedom ??

EU grant (via Foundation for Polish Science)

## **Jagellonian University International PhD Studies on Physics of Complex Systems**

- 1 M Euro
- 4 years
- 14 PhD students (1/2 - 2 years abroad)
- 17 Foreign Partners: J. Ambjorn, H. Nicolai, S. Sharpe, J.P. Blaizot ...
- 8 Local Supervisors