Towards a data-driven analysis of hadronic light-by-light scattering in the anomalous magnetic moment of the muon

# Martin Hoferichter



INSTITUTE for NUCLEAR THEORY Institute for Nuclear Theory University of Washington



KITP program on

## Frontiers in Nuclear Physics

#### Santa Barbara, October 21, 2016

G. Colangelo, MH, M. Procura, P. Stoffer, JHEP 09 (2014) 091, JHEP 09 (2015) 074

G. Colangelo, MH, B. Kubis, M. Procura, P. Stoffer, PLB 738 (2014) 6

MH, B. Kubis, S. Leupold, F. Niecknig, S. Schneider, EPJC 74 (2014) 3180

lots of work in progress

Anomalous magnetic moments a<sub>ℓ</sub>
 → prime low-energy precision observables

$$oldsymbol{a}_{oldsymbol{\ell}} = rac{g_{oldsymbol{\ell}}-2}{2} \qquad \mu = -grac{e}{2m}oldsymbol{S} \qquad \mathcal{H} = -\mu\cdotoldsymbol{B}$$

• Experimental precision 0.5 ppm BNL E821 2006

 $a_{\mu}^{\exp} = (116\,592\,089\pm63) \times 10^{-11}$ 

- Theory error of similar size
- Deviation from SM prediction around 3σ



#### • Experimental precision 0.5 ppm BNL E821 2006

 $a_{\mu}^{\exp} = (116592089 \pm 63) \times 10^{-11}$ 

- New experiment at **FNAL** (E989) aiming at 0.14 ppm, data taking to start in 2017  $\Rightarrow \Delta a_{\mu}^{exp} = 15 \times 10^{-11}$  as reference point
- J-PARC E821 statistics goal, new approach with ultra-cold muons, R&D in progress, EDM
- Comparison in review on "Precision muon physics" Gorringe, Hertzog 2015
  - $\Rightarrow$  Need to improve theory accordingly

1= nac

	$\frac{a_{\mu}}{2}$ [10 <sup>-11</sup> ]	$\Delta a_{\mu} [10^{-11}]$
experiment	116 592 089.	63.
$QED\ \mathcal{O}(\alpha)$	116 140 973.21	0.03
QED $\mathcal{O}(\alpha^2)$	413 217.63	0.01
QED $\mathcal{O}(\alpha^3)$	30 141.90	0.00
$QED\ \mathcal{O}(lpha^4)$	381.01	0.02
QED $\mathcal{O}(\alpha^5)$	5.09	0.01
QED total	116 584 718.85	0.04
electroweak, total	153.6	1.0
HVP (LO)	6 949.	43.
HVP (NLO)	-98.	1.
HLbL (LO)	116.	40.
HVP (NNLO)	12.4	0.1
HLbL (NLO)	3.	2.
theory	116 591 855.	59.



Schwinger 1948

= 200

	$a_{\mu}$ [10 <sup>-11</sup> ]	$\Delta a_{\mu} [10^{-11}]$
experiment	116 592 089.	63.
$QED\ \mathcal{O}(lpha)$	116 140 973.21	0.03
QED $\mathcal{O}(\alpha^2)$	413 217.63	0.01
QED $\mathcal{O}(\alpha^3)$	30 141.90	0.00
QED $\mathcal{O}(\alpha^4)$	381.01	0.02
QED $\mathcal{O}(\alpha^5)$	5.09	0.01
QED total	116 584 718.85	0.04
electroweak, total	153.6	1.0
HVP (LO)	6 949.	43.
HVP (NLO)	-98.	1.
HLbL (LO)	116.	40.
HVP (NNLO)	12.4	0.1
HLbL (NLO)	3.	2.
theory	116 591 855.	59.



Sommerfield, Petermann 1957

-

	$a_{\mu}$ [10 <sup>-11</sup> ]	$\Delta a_{\mu} [10^{-11}]$
experiment	116 592 089.	63.
$QED\ \mathcal{O}(lpha)$	116 140 973.21	0.03
QED $O(\alpha^2)$	413 217.63	0.01
QED $\mathcal{O}(\alpha^3)$	30 141.90	0.00
$QED\ \mathcal{O}(lpha^4)$	381.01	0.02
QED $\mathcal{O}(\alpha^5)$	5.09	0.01
QED total	116 584 718.85	0.04
electroweak, total	153.6	1.0
HVP (LO)	6 949.	43.
HVP (NLO)	-98.	1.
HLbL (LO)	116.	40.
HVP (NNLO)	12.4	0.1
HLbL (NLO)	3.	2.
theory	116 591 855.	59.



Kinoshita et al. 2012

M. Hoferichter (Institute for Nuclear Theory)

-

	$a_{\mu}$ [10 <sup>-11</sup> ]	$\Delta a_{\mu} [10^{-11}]$
experiment	116 592 089.	63.
$QED\ \mathcal{O}(lpha)$	116 140 973.21	0.03
QED $O(\alpha^2)$	413 217.63	0.01
QED $\mathcal{O}(\alpha^3)$	30 141.90	0.00
$QED\ \mathcal{O}(lpha^4)$	381.01	0.02
QED $\mathcal{O}(\alpha^5)$	5.09	0.01
QED total	116 584 718.85	0.04
electroweak, total	153.6	1.0
HVP (LO)	6 949.	43.
HVP (NLO)	-98.	1.
HLbL (LO)	116.	40.
HVP (NNLO)	12.4	0.1
HLbL (NLO)	3.	2.
theory	116 591 855.	59.



1-loop: Jackiw, Weinberg and others 1972 2-loop: Kukhto et al. 1992, Czarnecki, Krause, Marciano 1995, Degrassi, Giudice 1998, Knecht, Peris, Perrottet, de Rafael 2002, Vainshtein 2003, Heinemeyer, Stöckinger, Weiglein 2004, Gribouk, Czarnecki 2005 Update after Higgs discovery: Gnendiger et al. 2013

M. Hoferichter (Institute for Nuclear Theory) Towards a data-driven

	$a_{\mu}$ [10 <sup>-11</sup> ]	$\Delta a_{\mu} [10^{-11}]$
experiment	116 592 089.	63.
$QED\ \mathcal{O}(lpha)$	116 140 973.21	0.03
QED $\mathcal{O}(\alpha^2)$	413 217.63	0.01
QED $\mathcal{O}(\alpha^3)$	30 141.90	0.00
QED $\mathcal{O}(\alpha^4)$	381.01	0.02
QED $\mathcal{O}(\alpha^5)$	5.09	0.01
QED total	116 584 718.85	0.04
electroweak, total	153.6	1.0
HVP (LO)	6 949.	43.
HVP (NLO)	-98.	1.
HLbL (LO)	116.	40.
HVP (NNLO)	12.4	0.1
HLbL (NLO)	3.	2.
theory	116 591 855.	59.



Hagiwara et al. 2011

ъ

	$a_{\mu}$ [10 <sup>-11</sup> ]	$\Delta a_{\mu} [10^{-11}]$
experiment	116 592 089.	63.
$QED\ \mathcal{O}(lpha)$	116 140 973.21	0.03
QED $O(\alpha^2)$	413 217.63	0.01
QED $\mathcal{O}(\alpha^3)$	30 141.90	0.00
QED $\mathcal{O}(\alpha^4)$	381.01	0.02
QED $\mathcal{O}(\alpha^5)$	5.09	0.01
QED total	116 584 718.85	0.04
electroweak, total	153.6	1.0
HVP (LO)	6 949.	43.
HVP (NLO)	-98.	1.
HLbL (LO)	116.	40.
HVP (NNLO)	12.4	0.1
HLbL (NLO)	3.	2.
theory	116 591 855.	59.



Calmet et al. 1976, Hagiwara et al. 2011

-

	$a_{\mu}$ [10 <sup>-11</sup> ]	$\Delta a_{\mu} [10^{-11}]$
experiment	116 592 089.	63.
$QED\ \mathcal{O}(lpha)$	116 140 973.21	0.03
QED $\mathcal{O}(\alpha^2)$	413 217.63	0.01
QED $\mathcal{O}(\alpha^3)$	30 141.90	0.00
$QED\ \mathcal{O}(lpha^4)$	381.01	0.02
QED $\mathcal{O}(\alpha^5)$	5.09	0.01
QED total	116 584 718.85	0.04
electroweak, total	153.6	1.0
HVP (LO)	6 949.	43.
HVP (NLO)	-98.	1.
HLbL (LO)	116.	40.
HVP (NNLO)	12.4	0.1
HLbL (NLO)	3.	2.
theory	116 591 855.	59.



Hayakawa, Kinoshita, Sanda 1995 Bijnens, Pallante, Prades 1995 Knecht, Nyffeler 2001 Jegerlehner, Nyffeler 2009

M. Hoferichter (Institute for Nuclear Theory)

	$a_{\mu}$ [10 <sup>-11</sup> ]	$\Delta a_{\mu} [10^{-11}]$
experiment	116 592 089.	63.
$QED\ \mathcal{O}(lpha)$	116 140 973.21	0.03
QED $O(\alpha^2)$	413 217.63	0.01
QED $\mathcal{O}(\alpha^3)$	30 141.90	0.00
QED $\mathcal{O}(\alpha^4)$	381.01	0.02
QED $\mathcal{O}(\alpha^5)$	5.09	0.01
QED total	116 584 718.85	0.04
electroweak, total	153.6	1.0
HVP (LO)	6 949.	43.
HVP (NLO)	-98.	1.
HLbL (LO)	116.	40.
HVP (NNLO)	12.4	0.1
HLbL (NLO)	3.	2.
theory	116 591 855.	59.



Kurz, Liu, Marquard, Steinhauser 2014

	$a_{\mu}$ [10 <sup>-11</sup> ]	$\Delta a_{\mu} [10^{-11}]$
experiment	116 592 089.	63.
$QED\ \mathcal{O}(lpha)$	116 140 973.21	0.03
QED $\mathcal{O}(\alpha^2)$	413 217.63	0.01
QED $\mathcal{O}(\alpha^3)$	30 141.90	0.00
$QED\ \mathcal{O}(lpha^4)$	381.01	0.02
QED $\mathcal{O}(\alpha^5)$	5.09	0.01
QED total	116 584 718.85	0.04
electroweak, total	153.6	1.0
HVP (LO)	6 949.	43.
HVP (NLO)	-98.	1.
HLbL (LO)	116.	40.
HVP (NNLO)	12.4	0.1
HLbL (NLO)	3.	2.
theory	116 591 855.	59.



Colangelo, MH, Nyffeler, Passera, Stoffer 2014

4

	$a_{\mu}$ [10 <sup>-11</sup> ]	$\Delta a_{\mu} [10^{-11}]$
experiment	116 592 089.	63.
$QED\ \mathcal{O}(lpha)$	116 140 973.21	0.03
QED $\mathcal{O}(\alpha^2)$	413 217.63	0.01
QED $\mathcal{O}(\alpha^3)$	30 141.90	0.00
$QED\ \mathcal{O}(lpha^4)$	381.01	0.02
QED $\mathcal{O}(\alpha^5)$	5.09	0.01
QED total	116 584 718.85	0.04
electroweak, total	153.6	1.0
HVP (LO)	6 949.	43.
HVP (NLO)	-98.	1.
HLbL (LO)	116.	40.
HVP (NNLO)	12.4	0.1
HLbL (NLO)	3.	2.
theory	116 591 855.	59.

$$a_{\mu}^{\mathsf{exp}} - a_{\mu}^{\mathsf{SM}} = (234 \pm 86) imes 10^{-11} [2.7\sigma]$$

⇒Theory error comes almost exclusively from hadronic part

M. Hoferichter (Institute for Nuclear Theory) Towards a data-driven analysis of HLbL scattering

# Outline

## Hadronic vacuum polarization

- Approaches to HLbL
- The HLbL tensor: gauge invariance and crossing symmetry

## A dispersion relation for HLbL

- Master formula
- Pion box
- Pion rescattering
- Input for pion pole (and beyond)

## Summary and outlook

# Hadronic vacuum polarization

- General principles yield direct connection with experiment
  - Gauge invariance

$$k, \mu \qquad k, \nu \qquad = -i(k^2 g^{\mu\nu} - k^{\mu} k^{\nu}) \Pi(k^2)$$

Analyticity

$$\Pi_{\text{ren}} = \Pi(k^2) - \Pi(0) = \frac{k^2}{\pi} \int_{4M_\pi^2}^{\infty} \text{d}s \frac{\text{Im}\,\Pi(s)}{s(s-k^2)}$$

Unitarity

$$\operatorname{Im}\Pi(s) = \frac{s}{4\pi\alpha}\sigma_{\operatorname{tot}}(e^+e^- \to \operatorname{hadrons}) = \frac{\alpha}{3}R(s)$$

- 1 Lorentz structure, 1 kinematic variable, parameter-free
- Dedicated e<sup>+</sup>e<sup>-</sup> program under way: BaBar, Belle, BESIII, CMD3, KLOE2, SND (still hard to go much below 1%)

= ∕) < (~

## Hadronic vacuum polarization: two-pion channel

- Accuracy goal: 0.6% (present)  $\rightarrow$  0.2% (experiment)
- Systematics of  $\pi\pi$  channel:  $\tau$  data, ISR data
- Current status BESIII 2015





## HLbL: irreducible uncertainty?

$\pi^0, \eta, \eta'$	$\pi^{\pm}, K^{\pm}$		$\sigma, f_0,$		2	quarks	R
Contribution	BPP	HKS	KN	MV	BP	PdRV	N/JN
$r^0, \eta, \eta'$	$85 \pm 13$	$82.7 \pm 6.4$	$83 \pm 12$	$114 \pm 10$	-	$114 \pm 13$	$99 \pm 16$
r, K loops	$-19 \pm 13$	$-4.5 \pm 8.1$	-	-	-	$-19 \pm 19$	$-19 \pm 13$
$r, K$ loops + other subleading in $N_c$	-	-	-	$0 \pm 10$	-	-	-
xial vectors	$2.5 \pm 1.0$	$1.7 \pm 1.7$	-	$22 \pm 5$	-	$15 \pm 10$	$22 \pm 5$
calars	$-6.8 \pm 2.0$	-	-	-	-	$-7 \pm 7$	$-7 \pm 2$
Juark loops	$21\pm3$	$9.7\pm11.1$	-	-	-	2.3±	$21\pm3$
otal	$83 \pm 32$	$89.6\pm15.4$	$80\pm40$	$136\pm25$	$110 \pm 40$	$105\pm26$	$116 \pm 39$

• HVP systematically improvable

Jegerlehner, Nyffeler 2009

- HLbL more challenging
  - 4-point function of EM currents
  - Unambiguous definitions?
  - 5 kinematic variables, many more Lorentz structures (but only 7 master structures)
  - Folk theorem: "it cannot be expressed in terms of measurable quantities"
- Our suggestion: adapt methods from HVP, stay as data-driven as possible

# Approaches to HLbL

#### Model calculations

• ENJL	Bijnens, Pallante, Prades 1995-96
<ul> <li>NJL and hidden gauge</li> </ul>	Hayakawa, Kinoshita, Sanda 1995-96
<ul> <li>Nonlocal <i>χ</i>QM</li> </ul>	Dorokhov, Broniowski 2008
AdS/CFT	Cappiello, Cata, D'Ambrosio 2010
<ul> <li>Dyson–Schwinger</li> </ul>	Goecke, Fischer, Williams 2011
• Constituent $\chi$ QM	Greynat, de Rafael 2012
Resonances in narrow-width limit	Pauk, Vanderhaeghen 2014

### Rigorous constraints from QCD

<ul> <li>High-energy constraints ta</li> </ul>	ken into account in several	models above,
addressed specifically by		Knecht, Nyffeler 2001
• ChPT for $a_{\mu}$	Knecht, Nyffeler, Perrottet	, de Rafael 2002, Ramsey-Musolf, Wise 2002
<ul> <li>High-energy constraints related to the axial anomaly</li> </ul>		Melnikov, Vainshtein 2004, Nyffeler 2009
• Sum rules for $\gamma^*\gamma \to X$		Pascalutsa, Pauk, Vanderhaeghen 2012
<ul> <li>Low-energy constraints from pion polarizabilities</li> </ul>		Engel, Ramsey-Musolf 2013
Lattice		Blum et al. 2005, 2012-16, Green et al. 2015

A D b 4 A b

크 > < 크 >

1= 990

#### Cauchy's theorem

$$f(s) = rac{1}{2\pi i} \int_{\partial\Omega} rac{\mathrm{d}s' f(s')}{s' - s}$$



# From Cauchy's theorem to dispersion relations

#### Cauchy's theorem

$$f(s) = rac{1}{2\pi i} \int_{\partial\Omega} rac{\mathrm{d}s' f(s')}{s' - s}$$



## • Dispersion relation

$$f(s) = \frac{g}{s - M^2} + \frac{1}{\pi} \int_{\text{cuts}} \frac{\mathrm{d}s' \operatorname{Im} f(s')}{s' - s}$$

 $\hookrightarrow \textbf{analyticity}$ 



# From Cauchy's theorem to dispersion relations

#### • Dispersion relation

$$f(s) = \frac{g}{s - M^2} + \frac{1}{\pi} \int_{\text{cuts}} \frac{\mathrm{d}s' \operatorname{Im} f(s')}{s' - s}$$

 $\hookrightarrow \textbf{analyticity}$ 

Subtractions

$$f(s) = \frac{g}{s - M^2} + \underbrace{C}_{f(0) + \frac{g}{M^2}} + \frac{s}{\pi} \int_{\text{cuts}} \frac{\text{d}s' \,\text{Im} \, f(s')}{s'(s' - s)}$$



## Dispersion relation

$$f(s) = \frac{g}{s - M^2} + \frac{1}{\pi} \int_{\text{cuts}} \frac{\mathrm{d}s' \operatorname{Im} f(s')}{s' - s}$$

 $\hookrightarrow \textbf{analyticity}$ 

Subtractions

$$f(s) = \frac{g}{s - M^2} + \underbrace{C}_{f(0) + \frac{g}{M^2}} + \frac{s}{\pi} \int_{\text{cuts}} \frac{\text{d}s' \, \text{Im} \, f(s')}{s'(s' - s)}$$

Imaginary part from Cutkosky rules

 $\hookrightarrow \text{ forward direction: } \textbf{optical theorem}$ 

see HVP and  $\sigma(e^+e^- \rightarrow \text{hadrons})$ 

- Unitarity for partial waves:  $\lim f(s) = \rho(s)|f(s)|^2$
- Residue g reaction-independent



# Why dispersive approach?

- Analytic structure: poles and cuts
  - $\hookrightarrow$  **Residues** and **imaginary parts**  $\Rightarrow$  by definition **on-shell** quantities
  - $\hookrightarrow$  form factors and scattering amplitudes from experiment
  - ← model-independent definition of all contributions!

# Why dispersive approach?

- Analytic structure: poles and cuts
  - $\hookrightarrow$  **Residues** and **imaginary parts**  $\Rightarrow$  by definition **on-shell** quantities
  - $\hookrightarrow$  form factors and scattering amplitudes from experiment
  - ← model-independent definition of all contributions!
- Challenges
  - Find suitable quantities for dispersive analysis: Bardeen–Tung–Tarrach basis
  - Large number of amplitudes and invariants: no closed formula as for HVP
    - $\hookrightarrow$  Expansion in mass of intermediate states and partial waves
- Pseudoscalar poles most important, next  $\pi\pi$  cuts
- Decompose the tensor according to

$$\Pi_{\mu\nu\lambda\sigma} = \Pi^{\pi^{0}\text{-pole}}_{\mu\nu\lambda\sigma} + \Pi^{\pi\text{-box}}_{\mu\nu\lambda\sigma} + \bar{\Pi}_{\mu\nu\lambda\sigma} + \cdots$$

- $\hookrightarrow$  accounts for **one-** and **two-pion** intermediate states
- Generalizes immediately to  $\eta$ ,  $\eta'$ ,  $K\bar{K}$ , but e.g.  $3\pi$  more difficult

$$q_4 = k = q_1 + q_2 + q_3$$
  $k^2 = 0$ 



HLbL tensor

$$\Pi^{\mu\nu\lambda\sigma} = i^{3} \int d^{4}x \int d^{4}y \int d^{4}z \, e^{-i(q_{1}\cdot x + q_{2}\cdot y + q_{3}\cdot z)} \langle 0|T\{j^{\mu}(x)j^{\nu}(y)j^{\lambda}(z)j^{\sigma}(0)\}|0\rangle$$
$$= g^{\mu\nu}g^{\lambda\sigma}\Pi_{1} + g^{\mu\lambda}g^{\nu\sigma}\Pi_{2} + g^{\mu\sigma}g^{\nu\lambda}\Pi_{3} + \sum_{ijkl} q^{\mu}_{i}q^{\nu}_{j}q^{\lambda}_{k}q^{\sigma}_{l}\Pi_{ijkl} + \cdots$$

- Lorentz decomposition: 138 (136 Eichmann, Fischer, Heupel, Williams 2014) functions
- Constraints from gauge invariance: Bardeen, Tung 1968, Tarrach 1975
- Need basis free of kinematic singularities and zeros

## Detour: subprocess $\gamma^* \gamma^* \to \pi \pi$

• Consider 
$$\gamma^*(q_1, \lambda_1)\gamma^*(q_2, \lambda_2) \to \pi(p_1)\pi(p_2)$$
  
 $W^{\mu\nu} = i \int d^4x \, e^{-iq_1 \cdot x} \langle \pi(p_1)\pi(p_2)|T\{j^{\mu}(x)j^{\nu}(0)|0\rangle$   
 $= g^{\mu\nu}W_1 + \sum_{ij} q_i^{\mu}q_j^{\nu}W_2^{ij} \qquad q_3 = p_2 - p_1$ 

- Lorentz decomposition: 10 scalar functions
- Gauge invariance:

$$q_1^{\mu} \mathbf{W}_{\mu\nu} = q_2^{\nu} \mathbf{W}_{\mu\nu} = 0$$

• Bardeen, Tung 1968: hit with projectors  $I^{\mu
u} = g^{\mu
u} - rac{q_2^{\mu}q_1^{\nu}}{q_1\cdot q_2}$ 

$$W_{\mu\nu} = I_{\mu\mu'} I_{\nu'\nu} W_{\mu'\nu'} = \sum_{i=1}^{5} \bar{T}^{i}_{\mu\nu} \bar{A}_{i} = \sum_{i=1}^{5} T^{i}_{\mu\nu} A_{i}$$

•  $\bar{A}_i$  free of kinematic singularities, but not zeros  $\hookrightarrow$  remove poles from  $\bar{T}_i^{\mu}$  to get to  $A_i$ 

#### The resulting basis

$$\begin{split} T_1^{\mu\nu} &= q_1 \cdot q_2 g^{\mu\nu} - q_2^{\mu} q_1^{\nu} \\ T_2^{\mu\nu} &= q_1^2 q_2^2 g^{\mu\nu} + q_1 \cdot q_2 q_1^{\mu} q_2^{\nu} - q_1^2 q_2^{\mu} q_2^{\nu} - q_2^2 q_1^{\mu} q_1^{\nu} \\ T_3^{\mu\nu} &= q_1^2 q_2 \cdot q_3 g^{\mu\nu} + q_1 \cdot q_2 q_1^{\mu} q_3^{\nu} - q_1^2 q_2^{\mu} q_2^{\nu} - q_2^2 q_1^{\mu} q_1^{\nu} \\ T_4^{\mu\nu} &= q_2^2 q_1 \cdot q_3 g^{\mu\nu} + q_1 \cdot q_2 q_3^{\mu} q_2^{\nu} - q_2^2 q_3^{\mu} q_1^{\nu} - q_1 \cdot q_3 q_2^{\mu} q_2^{\nu} \\ T_5^{\mu\nu} &= q_1 \cdot q_3 q_2 \cdot q_3 g^{\mu\nu} + q_1 \cdot q_2 q_3^{\mu} q_3^{\nu} - q_1 \cdot q_3 q_2^{\mu} q_3^{\nu} - q_2 \cdot q_3 q_1^{\mu} q_1^{\nu} \end{split}$$

becomes degenerate for  $q_1 \cdot q_2 = 0$  Tarrach 1975

Need one more structure

$$T_{6}^{\mu\nu} = \left(q_{1}^{2}q_{3}^{\mu} - q_{1} \cdot q_{3}q_{1}^{\mu}\right)\left(q_{2}^{2}q_{3}^{\nu} - q_{2} \cdot q_{3}q_{2}^{\nu}\right)$$

#### $\hookrightarrow$ redundant set of 5 + 1 BTT functions

Crossing symmetry of the pions actually removes Tarrach ambiguity Drechsel et al. 1998

= 990

- BTT for HLbL Colangelo, MH, Procura, Stoffer 2015
  - 43 basis tensors
  - I1 additional ones
  - Out of 54 only 7 independent (up to crossing)
  - 2 further redundancies in *d* = 4



$$\Pi^{\mu\nu\lambda\sigma} = \sum_{i=1}^{54} T_i^{\mu\nu\lambda\sigma} \Pi_i$$

## Back to HLbL

- BTT for HLbL Colangelo, MH, Procura, Stoffer 2015
  - 43 basis tensors
  - I1 additional ones
  - Out of 54 only 7 independent (up to crossing)
  - 2 further redundancies in *d* = 4



$$\begin{split} T_{1}^{\mu\nu\lambda\sigma} &= \epsilon^{\mu\nu\alpha\beta} \epsilon^{\lambda\sigma\gamma\delta} a_{1\,\alpha} q_{2\,\beta} q_{3\,\gamma} q_{4\,\delta} \qquad T_{4}^{\mu\nu\lambda\sigma} = \left(q_{2}^{\mu} q_{1}^{\nu} - q_{1} \cdot q_{2} g^{\mu\nu}\right) \left(q_{4}^{\lambda} q_{3}^{\sigma} - q_{3} \cdot q_{4} g^{\lambda\sigma}\right) \\ T_{7}^{\mu\nu\lambda\sigma} &= \left(q_{2}^{\mu} q_{1}^{\nu} - q_{1} \cdot q_{2} g^{\mu\nu}\right) \left(q_{1} \cdot q_{4} \left(q_{1}^{\lambda} q_{3}^{\sigma} - q_{1} \cdot q_{3} g^{\lambda\sigma}\right) + q_{4}^{\lambda} q_{1}^{\sigma} q_{1} \cdot q_{3} - q_{1}^{\lambda} q_{1}^{\sigma} q_{3} \cdot q_{4}\right) \\ T_{19}^{\mu\nu\lambda\sigma} &= \left(q_{2}^{\mu} q_{1}^{\nu} - q_{1} \cdot q_{2} g^{\mu\nu}\right) \left(q_{2} \cdot q_{4} \left(q_{1}^{\lambda} q_{3}^{\sigma} - q_{1} \cdot q_{3} g^{\lambda\sigma}\right) + q_{4}^{\lambda} q_{2}^{\sigma} q_{1} \cdot q_{3} - q_{1}^{\lambda} q_{2}^{\sigma} q_{3} \cdot q_{4}\right) \\ T_{31}^{\mu\nu\lambda\sigma} &= \left(q_{2}^{\mu} q_{1}^{\nu} - q_{1} \cdot q_{2} g^{\mu\nu}\right) \left(q_{2}^{\lambda} q_{1} \cdot q_{3} - q_{1}^{\lambda} q_{2} \cdot q_{3}\right) \left(q_{2}^{\sigma} q_{1} \cdot q_{4} - q_{1}^{\sigma} q_{2} \cdot q_{4}\right) \\ &+ g^{\mu\sigma} \left(q_{2}^{\lambda} q_{3} \cdot q_{4} - q_{4}^{\mu} q_{1} \cdot q_{3}\right) \left(q_{3}^{\nu} q_{4}^{\lambda} q_{2}^{\sigma} - q_{4}^{\nu} q_{2}^{\lambda} q_{3}^{\sigma} + g^{\lambda\sigma} \left(q_{4}^{\nu} q_{2} \cdot q_{3} - q_{3}^{\nu} q_{2}^{\lambda} q_{4}\right) \\ &+ g^{\nu\sigma} \left(q_{2}^{\lambda} q_{3} \cdot q_{4} - q_{4}^{\lambda} q_{2} \cdot q_{3}\right) + g^{\lambda\nu} \left(q_{3}^{\sigma} q_{2} \cdot q_{4} - q_{2}^{\sigma} q_{3} \cdot q_{4}\right) \right) \\ T_{49}^{\mu\nu\lambda\sigma} &= q_{3}^{\sigma} \left(q_{1} \cdot q_{3} q_{2} \cdot q_{4} q_{4}^{\mu} q_{3}^{\lambda} q_{1}^{\lambda} + q_{1} \cdot q_{4} q_{2}^{\nu} q_{4}^{\lambda} q_{2}^{\lambda} q_{4} - q_{2}^{\sigma} q_{3} \cdot q_{4}\right) \right) \\ &+ q_{1} \cdot q_{4} q_{3}^{\mu} q_{4}^{\mu} q_{2}^{\lambda} q_{2}^{\lambda} - q_{2} \cdot q_{4} q_{4}^{\mu} q_{3}^{\lambda} q_{1}^{\lambda} + q_{1} \cdot q_{4} q_{2}^{\nu} q_{4} \left(q_{3}^{\mu} g^{\lambda\mu} - q_{3}^{\mu} g^{\lambda\nu}\right) \right) \\ &- q_{4}^{\lambda} \left(q_{1} \cdot q_{4} q_{2} \cdot q_{3} q_{3}^{\mu} g^{\mu\sigma} - q_{2} \cdot q_{3} q_{3}^{\mu} g^{\mu\sigma} + q_{1} \cdot q_{3} q_{2}^{\nu} q_{3} \left(q_{4}^{\nu} g^{\mu\sigma} - q_{4}^{\mu} g^{\nu\sigma}\right) \right) \\ &+ q_{1} \cdot q_{3} q_{4}^{\mu} q_{3}^{\mu} q_{3}^{\sigma} q_{2}^{\sigma} - q_{2} \cdot q_{3} q_{3}^{\mu} q_{3}^{\sigma} q_{1}^{\sigma} + q_{1} \cdot q_{3} q_{2}^{\mu} q_{3}^{\nu}\right) \left(q_{3}^{\mu} q_{1}^{\sigma} - q_{1} \cdot q_{3} g^{\mu\sigma}\right) \right) \\ &+ q_{1} \cdot q_{3} q_{4}^{\mu} q_{3}^{\mu} q_{3}^{\sigma} q_{1}^{\sigma} - q_{2} \cdot q_{3} q_{3}^{\mu} q_{1}^{\sigma}\right) \right) \\ &+ q_{1} \cdot q_{3} q_{4}^{\mu} q_{3}^{\mu} q_{3}^{\sigma} q_{1}^{\sigma} - q_{2} \cdot q_{3} q_{3}^{\mu}\right) \left(q_{3}^{\mu} q_{2}^{\sigma} - q_{2} \cdot q_{3} q_{3}^{\mu} q_{1}^{\sigma}$$

- BTT for HLbL Colangelo, MH, Procura, Stoffer 2015
  - 43 basis tensors
  - 11 additional ones
  - Out of 54 only 7 independent (up to crossing)
  - 2 further redundancies in *d* = 4



$$\Pi^{\mu\nu\lambda\sigma} = \sum_{i=1}^{54} T_i^{\mu\nu\lambda\sigma} \Pi_i$$

 $\hookrightarrow$  dynamical calculation for only 7 scalar amplitudes!

Master formula for  $a_{\mu}$ 

$$\mathbf{a}_{\mu}^{\mathsf{HLbL}} = -e^{6} \int \frac{\mathsf{d}^{4} q_{1}}{(2\pi)^{4}} \int \frac{\mathsf{d}^{4} q_{2}}{(2\pi)^{4}} \frac{\sum_{i=1}^{12} \hat{T}_{i}(q_{1}, q_{2}; p) \bar{\Pi}_{i}(q_{1}, q_{2}, -q_{1} - q_{2})}{q_{1}^{2} q_{2}^{2} (q_{1} + q_{2})^{2} ((p + q_{1})^{2} - m_{\mu}^{2}) ((p - q_{2})^{2} - m_{\mu}^{2})}$$

- $\hat{T}_i$ : known kernel functions
- $\overline{\Pi}_i$ : linear combinations of  $\Pi_i$
- Can perform five integrations with Gegenbauer polynomials

### Master formula for $a_{\mu}$

$$\mathbf{a}_{\mu}^{\text{HLbL}} = \frac{2\alpha^3}{3\pi^2} \int_0^\infty \mathrm{d}Q_1 \int_0^\infty \mathrm{d}Q_2 \int_{-1}^1 \mathrm{d}\tau \sqrt{1 - \tau^2} Q_1^3 Q_2^3 \sum_{i=1}^{12} T_i(Q_1, Q_2, \tau) \bar{\Pi}_i(Q_1, Q_2, \tau)$$

- T<sub>i</sub>: known kernel functions
- $\overline{\Pi}_i$ : linear combinations of  $\Pi_i$
- Can perform five integrations with Gegenbauer polynomials
- Wick rotation: all input quantities at space-like kinematics

## Master formula for $a_{\mu}$

$$\mathbf{a}_{\mu}^{\mathsf{HLbL}} = \frac{2\alpha^3}{3\pi^2} \int_0^\infty \mathsf{d}Q_1 \int_0^\infty \mathsf{d}Q_2 \int_{-1}^1 \mathsf{d}\tau \sqrt{1 - \tau^2} Q_1^3 Q_2^3 \sum_{i=1}^{12} T_i(Q_1, Q_2, \tau) \bar{\Pi}_i(Q_1, Q_2, \tau)$$

- *T<sub>i</sub>*: known kernel functions
- $\overline{\Pi}_i$ : linear combinations of  $\Pi_i$
- Can perform five integrations with Gegenbauer polynomials
- Wick rotation: all input quantities at space-like kinematics
- Decomposition completely general, now dispersion relations for n
  i
- Alternative: dispersion relations for Pauli form factor F2(t) Pauk, Vanderhaeghen 2014
  - $a_{\mu}^{\text{HLbL}}$  from  $a_{\mu} = F_2(0)$
  - Do the 2-loop integral dispersively, known result for pseudoscalar pole reproduced
  - Large number of cuts for higher intermediate states

ELE DQC

## Setting up the dispersive calculation: pion pole



- Pion pole: known
- Projection onto BTT basis: done
- Master formula reproduces explicit expressions in the literature
- To be done: incorporation of pQCD constraints

$$\Pi_{\mu\nu\lambda\sigma} = \Pi^{\pi^{0}\text{-pole}}_{\mu\nu\lambda\sigma} + \Pi^{\pi\text{-box}}_{\mu\nu\lambda\sigma} + \bar{\Pi}_{\mu\nu\lambda\sigma} + \cdots$$

In JHEP 2014 paper



Separate contribution with two simultaneous cuts

- Analytic properties like the box diagram in sQED
- Triangle and bulb required by gauge invariance
- Multiplication with vector form factor  $F_{\pi}^{V}$  gives correct  $q^{2}$ -dependence  $\Rightarrow$  FsQED

Claim: **FsQED** is not an approximation  $\Pi_{\mu\nu\lambda\sigma}^{\pi\text{-box}} = \Pi_{\mu\nu\lambda\sigma}^{\text{FsQED}}$


Now with BTT basis

- Constructed a Mandelstam representation for  $\pi\pi$  intermediate states with pion-pole left-hand cut
- Checked explicitly that this agrees with FsQED

Proven: **FsQED** is not an approximation  $\Pi^{\pi\text{-box}}_{\mu\nu\lambda\sigma} = \Pi^{\text{FsQED}}_{\mu\nu\lambda\sigma}$ Uniquely defines the notion of a "pion loop"

$$\Pi_{\mu\nu\lambda\sigma} = \Pi^{\pi^{0}\text{-pole}}_{\mu\nu\lambda\sigma} + \Pi^{\pi\text{-box}}_{\mu\nu\lambda\sigma} + \bar{\Pi}_{\mu\nu\lambda\sigma} + \cdots$$

- Remainder  $\bar{\Pi}_{\mu\nu\lambda\sigma}$  has cuts only in one channel
- Physics:  $\pi\pi$  rescattering
- Calculated with a partial-wave expansion
- Similar for  $\eta$ ,  $\eta'$  poles and  $K\bar{K}$  intermediate states



$$\Pi_{i}^{\pi^{0}\text{-pole}}(s,t,u) = \frac{\rho_{i;s}}{s - M_{\pi}^{2}} + \frac{\rho_{i;t}}{t - M_{\pi}^{2}} + \frac{\rho_{i;u}}{u - M_{\pi}^{2}}$$

$$\rho_{i,s} = \delta_{i1}F_{\pi^{0}\gamma^{*}\gamma^{*}}(q_{1}^{2}, q_{2}^{2})F_{\pi^{0}\gamma^{*}\gamma^{*}}(q_{3}^{2}, q_{4}^{2})$$

$$\rho_{i,t} = \delta_{i2}F_{\pi^{0}\gamma^{*}\gamma^{*}}(q_{1}^{2}, q_{3}^{2})F_{\pi^{0}\gamma^{*}\gamma^{*}}(q_{2}^{2}, q_{4}^{2})$$

$$\rho_{i,u} = \delta_{i3}F_{\pi^{0}\gamma^{*}\gamma^{*}}(q_{1}^{2}, q_{4}^{2})F_{\pi^{0}\gamma^{*}\gamma^{*}}(q_{2}^{2}, q_{3}^{2})$$

- Crucial ingredient: pion transition form factor  $F_{\pi^0\gamma^*\gamma^*}(q_1^2, q_2^2)$
- $\bullet\,$  Dispersive approach: pion on-shell  $\rightarrow\,$  data input

### Pion box: projection onto BTT

Г

Very compact expressions in BTT basis

$$\begin{split} I_{l}^{\pi-\text{box}}(q_{1}^{2},q_{2}^{2},q_{3}^{2}) &= F_{\pi}^{V}(q_{1}^{2})F_{\pi}^{V}(q_{2}^{2})F_{\pi}^{V}(q_{3}^{2})\frac{1}{16\pi^{2}}\int_{0}^{1}dx\int_{0}^{1-x}dy\,I_{l}(x,y)\\ I_{1}(x,y) &= -\frac{2}{3}\frac{(1-2y)(1-2x-2y)(1-6x(1-x))}{\Delta^{2}}\qquad I_{7}(x,y) = -\frac{4}{3}\frac{(1-2x)^{2}(1-2y)^{2}y(1-y)}{\Delta^{3}}\\ I_{4}(x,y) &= -\frac{2}{3}\frac{(1-2x)(1+2x(1-3x(1-2y)-6y(1-y)))}{\Delta^{2}}\qquad \dots\\ \Delta &= M_{\pi}^{2} - xyq_{1}^{2} - x(1-x-y)q_{2}^{2} - y(1-x-y)q_{3}^{2} \end{split}$$

- Manifestly free of kinematic singularities
- Only 9 independent functions due to remaining crossing symmetries, e.g.

$$\Pi_{2} = \mathcal{C}_{23}[\Pi_{1}] \qquad \Pi_{5} = \mathcal{C}_{23}[\Pi_{4}] \qquad \Pi_{9} = \mathcal{C}_{13}[\mathcal{C}_{23}[\Pi_{7}]] \qquad \Pi_{10} = \mathcal{C}_{23}[\Pi_{7}]$$

#### and even just 6 independent $\overline{\Pi}_i$

### Pion box: numerics



- Only input space-like pion vector form factor
- Preliminary numbers:  $a_{\mu}^{\pi\text{-box}} = -15.9 \times 10^{-11}$ ,  $a_{\mu}^{\pi\text{-box,VMD}} = -16.4 \times 10^{-11}$
- Compare:  $a_{\mu}^{K\text{-box,VMD}} = -0.5 \times 10^{-11}$

Contribution	BPP	HKS	KN	MV	BP	PdRV	N/JN
$\pi^0, \eta, \eta'$	$85\pm13$	$82.7\pm6.4$	$83 \pm 12$	$114\pm10$	-	$114\pm13$	$99\pm16$
$\pi, K$ loops	$-19 \pm 13$	$-4.5 \pm 8.1$	-	-	-	$-19 \pm 19$	$-19 \pm 13$
$\pi$ , K loops + other subleading in $N_c$	-	-	-	$0 \pm 10$	-	-	-
Axial vectors	$2.5 \pm 1.0$	$1.7 \pm 1.7$	-	$22 \pm 5$	-	$15 \pm 10$	$22 \pm 5$
Scalars	$-6.8 \pm 2.0$	-	-	-	-	$-7 \pm 7$	$-7 \pm 2$
Quark loops	$21\pm3$	$9.7\pm11.1$	-	-	-	2.3±	$21\pm3$
Total	$83\pm32$	$89.6 \pm 15.4$	$80\pm40$	$136\pm25$	$110\pm40$	$105\pm26$	$116 \pm 39$

Only input space-like pion vector form factor

- Preliminary numbers:  $a_{\mu}^{\pi\text{-box}} = -15.9 \times 10^{-11}$ ,  $a_{\mu}^{\pi\text{-box},\text{VMD}} = -16.4 \times 10^{-11}$
- Compare:  $a_{\mu}^{K\text{-box,VMD}} = -0.5 \times 10^{-11}$

▲目▶ 目目 の々で

### Pion box: saturation



- Impose cutoff in momenta *Q*<sub>max</sub> (polar-coordinate-type trafo)
- Rapid convergence:  $Q_{\text{max}} = \{1, 1.5\} \text{ GeV} \Rightarrow a_{\mu}^{\pi\text{-box}} = \{95, 99\}\%$  of full result

• Dispersion relations for  $\Pi_i$ , e.g. fixed-*u* at  $u = u_b = q_1^2$ 

$$\Pi_{1}(q_{1}^{2}, q_{2}^{2}, q_{3}^{2}) = \lim_{q_{4}^{2} \to 0} \left( \frac{1}{\pi} \int_{4M_{\pi}^{2}}^{\infty} ds' \frac{D_{1}^{s;u}(s'; u_{b})}{s' - q_{3}^{2}} + \frac{1}{\pi} \int_{4M_{\pi}^{2}}^{\infty} dt' \frac{D_{1}^{t;u}(t'; u_{b})}{t' - q_{2}^{2}} \right)$$

• Discontinuities from unitarity: diagonal in helicity basis for partial waves, e.g.

$$\operatorname{Im} h_{++,++}^{J}(s; q_{1}^{2}, q_{2}^{2}; q_{3}^{2}, 0) = \frac{\sigma(s)}{16\pi} h_{J,++}^{*}(s; q_{1}^{2}, q_{2}^{2}) h_{J,++}(s; q_{3}^{2}, 0)$$



 $\hookrightarrow$  need to project onto BTT basis

Solved for S-waves in 2014, now for arbitrary partial waves

- BTT and d = 4 ambiguities lead to different representation for  $a_{\mu} \checkmark$
- Equivalence implies set of sum rules: checked with FsQED  $\checkmark$
- Projection on partial waves violates these sum rules if expansion is truncated
   → how fast is the convergence? ✓
- Unphysical photon polarizations seemed to contribute to  $a_{\mu} \checkmark$
- $\bullet\,$  Test case: partial-wave expansion of FsQED  $\checkmark\,$

## Partial-wave expansion of FsQED



### • fixed-s, -t, -u dispersion relation

$\hookrightarrow$	equivalent	due to	crossing	symmetry
-------------------	------------	--------	----------	----------

- Fixed-s special: no s-channel cut, cancellations for J = 0
- Can use sum rules to optimize partial-wave convergence
- Beyond FsQED

$$a_{\mu} = rac{1}{2} \left( a_{\mu}^{ ext{fixed-}s} + a_{\mu}^{ ext{fixed-}t} + a_{\mu}^{ ext{fixed-}u} 
ight) + ext{higher cuts}$$

M. Hoferichter (Institute for Nuclear Theory)

	J = 20	extrapolation
s	-16.40	-
t	-16.33	-16.40
и	-16.26	-16.40

 $a_{\mu}^{\pi\text{-box}}(J) \sim J^{-n}, n pprox rac{5}{2}$ 

# $\gamma^*\gamma^* \to \pi\pi$ partial waves

### Roy(-Steiner) equations = Dispersion relations + partial-wave expansion

### + crossing symmetry + unitarity + gauge invariance

• **On-shell case**  $\gamma\gamma 
ightarrow \pi\pi$  García-Martín, Moussallam 2010, MH,

Phillips, Schat 2011, partial-wave analysis Dai, Pennington 2014

- Singly-virtual  $\gamma^* \gamma \rightarrow \pi \pi$  Moussallam 2013
- Doubly-virtual  $\gamma^*\gamma^* o \pi\pi$ : anomalous thresholds

Colangelo, MH, Procura, Stoffer arXiv:1309.6877



# $\gamma^*\gamma^* \to \pi\pi$ partial waves

### Roy(-Steiner) equations = Dispersion relations + partial-wave expansion

### + crossing symmetry + unitarity + gauge invariance

- **On-shell case**  $\gamma\gamma \rightarrow \pi\pi$  García-Martín, Moussallam 2010, MH, Phillips, Schat 2011, **partial-wave analysis** Dai, Pennington 2014
- Singly-virtual  $\gamma^* \gamma \rightarrow \pi \pi$  Moussallam 2013
- Doubly-virtual  $\gamma^* \gamma^* \rightarrow \pi \pi$ : anomalous thresholds Colangelo, MH, Procura, Stoffer arXiv:1309.6877
- Constraints
  - Low energies: pion polarizabilities, ChPT
  - **Primakoff**:  $\gamma \pi \rightarrow \gamma \pi$  (COMPASS),  $\gamma \gamma \rightarrow \pi \pi$  (JLab)
  - Scattering:  $e^+e^- \rightarrow e^+e^-\pi\pi$ ,  $e^+e^- \rightarrow \pi\pi\gamma$
  - (Transition) Form factors:  $F_V^{\pi}$ ,  $\omega, \phi \to \pi^0 \gamma^*$





## Physics of $\gamma^*\gamma^* \to \pi\pi$

- ππ rescattering includes dofs corresponding to resonances, e.g. f<sub>2</sub>(1270)
- S-wave provides model-independent implementation of the  $f_0(500)$



## Physics of $\gamma^*\gamma^* \to \pi\pi$

- ππ rescattering includes dofs corresponding to resonances, e.g. f<sub>2</sub>(1270)
- S-wave provides model-independent implementation of the  $f_0(500)$
- Analytic continuation with dispersion theory: resonance properties
  - Precise determination of  $\sigma$ -pole parameters from  $\pi\pi$  scattering Caprini, Colangelo, Leutwyler 2006

$$M_{\sigma} = 441^{+16}_{-8} \,\mathrm{MeV} \qquad \Gamma_{\sigma} = 544^{+18}_{-25} \,\mathrm{MeV}$$

• Coupling  $\sigma \to \gamma \gamma$  from  $\gamma \gamma \to \pi \pi$  MH, Phillips, Schat 2011



糸(500) PARTIAL WIDTHS





M. Hoferichter (Institute for Nuclear Theory)

Towards a data-driven analysis of HLbL scattering

## Preliminary results for $\pi\pi$ rescattering

### Full analysis requires careful study of

- Subtractions
- Asymptotic behavior
- Structure of the left-hand cut
- Coupled-channel system of  $\pi\pi/\bar{K}K$
- Here: numerics for ππ rescattering with a pion-pole left-hand cut and phase shifts from inverse-amplitude method
  - Isolates  $\pi\pi$  states
  - Reproduces f<sub>0</sub>(500) properties and low-energy phenomenological phase shifts
  - Defines reasonable extrapolation to  $\infty$
  - Pion form factor still describes off-shell behavior
  - $\hookrightarrow$  solve dispersion relation for  $\gamma^*\gamma^* \to \pi\pi$  **S-waves**

### • S-wave contributions

cutoff	1 GeV	1.5 GeV	2 GeV	$\infty$
<i>l</i> = 0	-9.2	-9.5	-9.3	-8.8
<i>l</i> = 2	2.0	1.3	1.1	0.9

- Check on  $\gamma^*\gamma^* \to \pi\pi$ : sum rule involving J = 0 (and higher) amplitudes
  - $\hookrightarrow$  fulfilled at better than 10% with S-waves alone
- " $f_0(500)$  contribution" to  $a_{\mu}$  around  $-9 \times 10^{-11}$

## Preliminary results for $\pi\pi$ rescattering

Contribution	BPP	HKS	KN	MV	BP	PdRV	N/JN
$\pi^0, \eta, \eta'$	$85 \pm 13$	$82.7\pm6.4$	$83 \pm 12$	$114\pm10$	-	$114\pm13$	$99 \pm 16$
$\pi, K$ loops	$-19 \pm 13$	$-4.5 \pm 8.1$	-	-	-	$-19 \pm 19$	$-19 \pm 13$
$\pi$ , K loops + other subleading in N <sub>c</sub>	-	-	-	$0 \pm 10$	-	-	-
Axial vectors	$2.5 \pm 1.0$	$1.7 \pm 1.7$	-	$22 \pm 5$	-	$15 \pm 10$	$22 \pm 5$
Scalars	$-6.8 \pm 2.0$	-	-	-	-	$-7 \pm 7$	$-7 \pm 2$
Quark loops	$21\pm3$	$9.7 \pm 11.1$	-	-	-	2.3±	$21\pm3$
Total	$83\pm32$	$89.6\pm15.4$	$80\pm40$	$136\pm25$	$110\pm40$	$105\pm26$	$116 \pm 39$

### • S-wave contributions

cutoff	1 GeV	1.5 GeV	2 GeV	$\infty$
<i>l</i> = 0	-9.2	-9.5	-9.3	-8.8
<i>l</i> = 2	2.0	1.3	1.1	0.9

• Check on  $\gamma^* \gamma^* \to \pi \pi$ : sum rule involving J = 0 (and higher) amplitudes

- $\hookrightarrow$  fulfilled at better than 10% with S-waves alone
- " $f_0(500)$  contribution" to  $a_{\mu}$  around  $-9 \times 10^{-11}$

## Back to the pion pole: pion transition form factor

- In principle, the doubly-virtual form factor  $F_{\pi^0\gamma^*\gamma^*}(q_1^2,q_2^2)$  can be measured
- Absent data, and/or to improve accuracy: dispersive reconstruction
- Required input
  - Pion vector form factor
  - $\gamma^* \rightarrow 3\pi$  amplitude
  - $\pi\pi$  scattering amplitude
- Done for the singly-virtual case MH, Kubis, Leupold, Niecknig, Schneider 2014, doubly-virtual in progress
- Transition form factors  $\omega, \phi \to \pi^0 \gamma^*$  probe a particular doubly-virtual configuration



# Predicting $\sigma(e^+e^- \to \pi^0\gamma)$ from $\sigma(e^+e^- \to 3\pi)$



- Fit dispersive representation to
  - $e^+e^- 
    ightarrow 3\pi$
- Determines singly-virtual form factor in time-like region
- ③ Predict  $e^+e^- 
  ightarrow \pi^0\gamma$  as check on the

formalism

## Extraction of slope and space-like continuation

For HLbL need the form factor in the

### space-like region

 $\hookrightarrow$  another dispersion relation

$$\mathcal{F}_{\pi^0\gamma^*\gamma}(q^2,0) = \mathcal{F}_{\pi\gamma\gamma} + rac{q^2}{\pi} \int_{s_{ ext{thr}}}^{\infty} \mathrm{d}s' rac{\mathrm{Im}\, \mathcal{F}_{\pi^0\gamma^*\gamma}(s',0)}{s'(s'-q^2)}$$

• Sum rules for  $F_{\pi\gamma\gamma}$  and slope parameters

$$\begin{split} a_{\pi} &= \frac{M_{\pi^0}^2}{F_{\pi\gamma\gamma}} \frac{1}{\pi} \int_{s_{\text{thr}}}^{\infty} \mathrm{d}s' \frac{\mathrm{Im} \, F_{\pi^0\gamma^*\gamma}(s',0)}{s'^2} \\ &= (30.7 \pm 0.6) \times 10^{-3} \\ b_{\pi} &= (1.10 \pm 0.02) \times 10^{-3} \end{split}$$

- Soon to be tested at BESIII
- Similar program for  $\eta$ ,  $\eta'$ Hanhart, Kupść, Meißner, Stollenwerk, Wirzba 2013 Kubis, Plenter 2015, Xiao et al. 2015



## Left-hand cut



- **Pion pole**: coupling determined by  $F_V^{\pi}$  as before
- Multi-pion intermediate states: approximate in terms of resonances
  - $2\pi \sim \rho$ : can even be done **exactly** using  $\gamma^* \rightarrow 3\pi$  amplitude
    - ↔ cf. pion transition form factor MH, Kubis, Sakkas 2012, MH, Kubis, Leupold, Niecknig, Schneider 2014
  - $3\pi \sim \omega, \phi$ : narrow-width approximation
    - $\hookrightarrow$  transition form factors for  $\omega,\phi o \pi^0\gamma^*$  Schneider, Kubis, Niecknig 2012
  - Higher intermediate states also potentially relevant: axials, tensors
    - $\hookrightarrow$  sum rules to constrain their transition form factors Pauk, Vanderhaeghen 2014

## Towards a data-driven analysis of HLbL



- Reconstruction of  $\gamma^* \gamma^* \to \pi \pi, \pi^0$ : combine experiment and theory constraints
- Beyond:  $\eta$ ,  $\eta'$ ,  $K\bar{K}$ , multi-pion channels (resonances), pQCD constraints, ...

- Dispersive framework for the calculation of the HLbL contribution to a<sub>µ</sub>
- Includes one- and two-pion intermediate states, can be extended to other pseudoscalar poles and two-meson states
- General master formula in terms of BTT function
- Preliminary numbers for pion box and  $\pi\pi$  rescattering
- Next steps
  - Doubly-virtual pion transition form factor
  - Refined analysis of rescattering effects
  - Implementation of pQCD constraints
  - Error analysis: which input quantity has the biggest impact on  $a_{\mu}$ ?

## Pion transition form factor: physical regions



## Pion transition form factor: physical regions



M. Hoferichter (Institute for Nuclear Theory) Towards a data-driven a

Towards a data-driven analysis of HLbL scattering

Santa Barbara, October 21, 2016 38

### Pion transition form factor: physical regions



M. Hoferichter (Institute for Nuclear Theory)

Towards a data-driven analysis of HLbL scattering

Santa Barbara, October 21, 2016

### Unitarity for pion vector form factor

$$\operatorname{Im} F_V^{\pi}(s) = \theta(s - 4M_{\pi}^2) F_V^{\pi}(s) e^{-i\delta_1(s)} \sin \delta_1(s)$$

 $\hookrightarrow$  final-state theorem: phase of  $F_V^{\pi}$  equals  $\pi\pi P$ -wave phase  $\delta_1$  Watson 1954

### Unitarity for pion vector form factor

$$\operatorname{Im} F_V^{\pi}(s) = \theta(s - 4M_{\pi}^2) F_V^{\pi}(s) e^{-i\delta_1(s)} \sin \delta_1(s)$$

- $\hookrightarrow$  final-state theorem: phase of  $F_V^{\pi}$  equals  $\pi\pi$  *P*-wave phase  $\delta_1$  Watson 1954
- Solution in terms of Omnès function Omnès 1958

$$F_{V}^{\pi}(s) = P(s)\Omega_{1}(s) \qquad \Omega_{1}(s) = \exp\left\{\frac{s}{\pi}\int_{4M_{\pi}^{2}}^{\infty} ds' \frac{\delta_{1}(s')}{s'(s'-s)}\right\}$$

• Asymptotics + normalization  $\Rightarrow P(s) = 1$ 

### Unitarity

$$\operatorname{Im} f_1(s) = \theta(s - 4M_{\pi}^2) f_1(s) e^{-i\delta_1(s)} \sin \delta_1(s)$$



 $\hookrightarrow$  again Watson's theorem, but now left-hand cut in  $f_1(s)$ 

### Unitarity

$$\operatorname{Im} f_1(s) = \theta(s - 4M_{\pi}^2) f_1(s) e^{-i\delta_1(s)} \sin \delta_1(s)$$



 $\hookrightarrow$  again Watson's theorem, but now left-hand cut in  $f_1(s)$ 

Including the left-hand cut

$$\operatorname{Im} f_{1}(s) = \operatorname{Im} \mathcal{F}(s) = \left(\underbrace{\mathcal{F}(s)}_{\operatorname{RHC}} + \underbrace{\widehat{\mathcal{F}}(s)}_{\operatorname{LHC}}\right) \theta\left(s - 4M_{\pi}^{2}\right) \sin \delta_{1}(s) e^{-i\delta_{1}(s)}$$

 $f_1(s) = \mathcal{F}(s) + \hat{\mathcal{F}}(s) \qquad \hat{\mathcal{F}}(s) = 3\langle (1-z^2)\mathcal{F} \rangle \qquad \langle z^n \mathcal{F} \rangle = \frac{1}{2} \int_{-1}^1 dz \, z^n \mathcal{F}(t)$ 

Omnès solution for  $\mathcal{F}(s)$ 

$$\mathcal{F}(s) = \Omega_1(s) \left\{ \frac{C_1}{3} \left( 1 - \dot{\Omega}_1(0)s \right) + \frac{C_2}{3}s + \frac{s^2}{\pi} \int_{4M_{\pi}^2}^{\infty} ds' \frac{\hat{\mathcal{F}}(s')\sin\delta_1(s')}{s'^2(s'-s)|\Omega_1(s')|} \right\}$$

### Omnès solution for $\mathcal{F}(s)$

$$\mathcal{F}(s) = \Omega_1(s) \left\{ \frac{C_1}{3} \left( 1 - \dot{\Omega}_1(0)s \right) + \frac{C_2}{3}s + \frac{s^2}{\pi} \int_{4M_\pi^2}^{\infty} ds' \frac{\hat{\mathcal{F}}(s')\sin\delta_1(s')}{s'^2(s'-s)|\Omega_1(s')|} \right\}$$

Solve for *F*(s) by iteration

•  $\hat{\mathcal{F}}(s)$  corresponds to crossed-channel  $\pi\pi$  rescattering



• Important observation:  $\mathcal{F}(s)$  linear in  $C_i$ 

$$\mathcal{F}(s) = C_1 \mathcal{F}_1(s) + C_2 \mathcal{F}_2(s)$$

 $\hookrightarrow$  basis functions  $\mathcal{F}_i(s)$  can be calculated once and for all

## $\gamma\pi \to \pi\pi$ : from cross-section data to the transition form factor

- Representation of the cross section in terms of two parameters → fit C<sub>i</sub> to data мн, кubis, Sakkas 2012
  - Test of chiral anomaly  $F_{3\pi} = e/(4\pi^2 F_{\pi}^3)$
  - Precise description of f<sub>1</sub>
- Looking forward to COMPASS result
  - $\hookrightarrow$  currently: use chiral prediction



## $\gamma\pi \to \pi\pi$ : from cross-section data to the transition form factor

- Representation of the cross section in terms of two parameters → fit C<sub>i</sub> to data MH, Kubis, Sakkas 2012
  - Test of chiral anomaly  $F_{3\pi} = e/(4\pi^2 F_{\pi}^3)$
  - Precise description of f<sub>1</sub>
- Looking forward to COMPASS result
  - $\hookrightarrow$  currently: use chiral prediction
- Dispersion relation for  $f_{\pi^0\gamma}(s) = F_{\nu s}(s,0)$

$$f_{\pi^0\gamma}(s) = f_{\pi^0\gamma}(0) + \frac{s}{12\pi^2} \int_{4M_{\pi}^2}^{\infty} ds' \frac{q_{\pi}^3(s') (F_V^{\pi}(s'))^* f_1(s')}{s'^{3/2}(s'-s)}$$
$$q_{\pi}(s) = \sqrt{s/4 - M_{\pi}^2}$$





• Subtraction constant:  $f_{\pi^0\gamma}(0) = \frac{F_{\pi\gamma\gamma}}{2} = \frac{e^2}{8\pi^2 F_{\pi}}$ 

# $\omega, \phi \to \pi^0 \gamma^*$ transition form factor

- Similar procedure for  $\omega,\phi o 3\pi$  and  $\omega,\phi o \pi^0\gamma^*$  Schneider, Kubis, Niecknig 2012
- Additional complications due to decay kinematics



General virtualities: how to fix the normalization?

$$\hookrightarrow$$
  $F_{3\pi}$  for  $\gamma \pi \rightarrow \pi \pi$ , widths for  $\omega, \phi \rightarrow 3\pi$ 

• Fit to  $e^+e^- \rightarrow 3\pi$ 

$$\begin{aligned} \mathbf{a}(q^2) &= \alpha + \beta q^2 + \frac{q^4}{\pi} \int_{s_{\text{thr}}}^{\infty} \mathrm{d}s' \frac{\mathrm{Im}\,\mathcal{A}(s')}{s'^2(s'-q^2)} \\ \mathcal{A}(q^2) &= \frac{c_{\omega}}{M_{\omega}^2 - q^2 - i\sqrt{q^2}\Gamma_{\omega}(q^2)} + \frac{c_{\phi}}{M_{\phi}^2 - q^2 - i\sqrt{q^2}\Gamma_{\phi}(q^2)} \end{aligned}$$

•  $\alpha$  fixed by  $F_{3\pi}$ ,  $\Gamma_{\omega/\phi}(q^2)$  include  $3\pi$ ,  $K\bar{K}$ ,  $\pi^0\gamma$  channels

- Good analytic properties, free parameters:  $\beta$ ,  $c_{\omega}$ ,  $c_{\phi}$
- Valid up to 1.1 GeV, also fit including  $\omega'$ ,  $\omega''$  to estimate uncertainties

# Pion transition form factor: unitarity relations

process	unitarity relations	SC 1	SC 2	
	7: ~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~		$F_{\pi^0\gamma\gamma}$	$\gamma \pi \rightarrow \pi \pi$
	P	$F_{3\pi}$	$\sigma(\gamma\pi  o \pi\pi)$	
$\overset{\gamma^*_s}{\longrightarrow} \overset{\omega^*_{\gamma^*_s}}{\underset{i}{\longrightarrow}} \overset{\omega^*_{\gamma^*_s}}{\bigcup_{\gamma^*_s}} {\longrightarrow} {\overset}{\overset}{\longrightarrow} {\longrightarrow} {\overset}{\overset}{\overset}{\overset}{\overset}{\overset}{\overset}{\overset}{\overset}{\overset}{\overset}{\overset}{$	$\underbrace{\omega,\phi}_{i} \underbrace{ \begin{pmatrix} i \\ i \\ j \end{pmatrix}}_{i} \gamma_{v}^{*}$		$\Gamma_{\pi^0\gamma}$	$\omega \rightarrow 3\pi, \phi \rightarrow 3\pi$
		Γ <sub>3π</sub>	$rac{\mathrm{d}^2\Gamma}{\mathrm{d}s\mathrm{d}t}(\omega,\phi ightarrow 3\pi)$	
$\overset{\gamma^*_s}{\leadsto} {\bigvee} {\bigvee} {\bigvee} {\bigvee} {\underset{\iota_{\iota_{\iota_{\iota}}}}}$	7°		$\sigma(e^+e^-  o \pi^0\gamma)$	$\gamma^*  o {f 3}\pi$
		$\sigma(e^+e^-  ightarrow 3\pi)$	$\sigma(\gamma\pi o\pi\pi)$ $rac{\mathrm{d}^{2}\Gamma}{\mathrm{d}s\mathrm{d}t}(\omega,\phi o3\pi)$	resummation of
	Ϋ́,	$F_{3\pi}$	$\sigma(e^+e^-  ightarrow 3\pi)$	$\pi\pi$ rescattering

M. Hoferichter (Institute for Nuclear Theory)
# $\gamma^* \gamma^* \to \pi \pi$ partial waves: unitarity relations



M. Hoferichter (Institute for Nuclear Theory)

Towards a data-driven analysis of HLbL scattering

= 990

$$\bar{\Pi}^{\mu\nu\lambda\sigma} = \sum_{i} \left( A^{\mu\nu\lambda\sigma}_{i,s} \Pi_{i}(s) + A^{\mu\nu\lambda\sigma}_{i,t} \Pi_{i}(t) + A^{\mu\nu\lambda\sigma}_{i,u} \Pi_{i}(u) \right)$$

- Need to choose  $A_i^{\mu\nu\lambda\sigma}$  so that  $\Pi_i$  are free of kinematic singularities
- General procedure for finding such a basis Bardeen, Tung 1968, Tarrach 1975
- Results in non-diagonal terms

$$\Pi_{1}(\boldsymbol{s}) = \frac{\boldsymbol{s} - q_{3}^{2}}{\pi} \int_{4M_{\pi}^{2}}^{\infty} \frac{\mathrm{d}\boldsymbol{s}'}{\boldsymbol{s}' - q_{3}^{2}} \left( \mathcal{K}_{1}(\boldsymbol{s}', \boldsymbol{s}) \mathrm{Im} \, \bar{h}_{++,++}^{0}(\boldsymbol{s}') + \frac{2\xi_{1}\xi_{2}}{\lambda(\boldsymbol{s}', q_{1}^{2}, q_{2}^{2})} \mathrm{Im} \, \bar{h}_{00,++}^{0}(\boldsymbol{s}') \right)$$

### Example: $\gamma^* \gamma^* \to \pi \pi$

- Similar analysis for  $\gamma^*\gamma^* o \pi\pi$ : Bardeen-Tung-Tarrach basis
  - $\hookrightarrow$  partial-wave dispersion relations (**Roy–Steiner equations**)
- Find similar non-diagonal kernels

= 990

#### Example: $\gamma^* \gamma^* \rightarrow \pi \pi$

• Similar analysis for  $\gamma^*\gamma^* o \pi\pi$ : Bardeen-Tung-Tarrach basis

 $\hookrightarrow$  partial-wave dispersion relations (**Roy–Steiner equations**)

- Find similar non-diagonal kernels
- Check within 1-loop ChPT

$$\begin{split} &\frac{1}{\pi} \int_{4M_{\pi}^2}^{\infty} \mathrm{d}t' \left\{ \left( \frac{1}{t'-t} - \frac{t'-q_1^2 - q_2^2}{\lambda(t',q_1^2,q_2^2)} \right) \mathrm{Im} \, h_1(t';q_1^2,q_2^2) + \frac{2q_1^2q_2^2}{\lambda(t',q_1^2,q_2^2)} \mathrm{Im} \, h_2(t';q_1^2,q_2^2) \right\} \\ &= 1 + 2 \left( M_{\pi}^2 + \frac{tq_1^2q_2^2}{\lambda(t,q_1^2,q_2^2)} \right) C_0(t,q_1^2,q_2^2) + \frac{t(q_1^2 + q_2^2) - (q_1^2 - q_2^2)^2}{\lambda(t,q_1^2,q_2^2)} \bar{J}(t) \\ &- \frac{q_1^2(t+q_2^2 - q_1^2)}{\lambda(t,q_1^2,q_2^2)} \bar{J}(q_1^2) - \frac{q_2^2(t+q_1^2 - q_2^2)}{\lambda(t,q_1^2,q_2^2)} \bar{J}(q_2^2) \\ \mathrm{Im} \, h_1(t;q_1^2,q_2^2) = 2 \left( M_{\pi}^2 + \frac{tq_1^2q_2^2}{\lambda(t,q_1^2,q_2^2)} \right) \mathrm{Im} \, C_0(t,q_1^2,q_2^2) + \frac{t(q_1^2 + q_2^2) - (q_1^2 - q_2^2)^2}{\lambda(t,q_1^2,q_2^2)} \mathrm{Im} \, \bar{J}(t) \\ \mathrm{Im} \, h_2(t;q_1^2,q_2^2) = -\frac{1}{\lambda(t,q_1^2,q_2^2)} \left[ \left( t^2 - (q_1^2 - q_2^2)^2 \right) \mathrm{Im} \, C_0(t,q_1^2,q_2^2) + 4t \mathrm{Im} \, \bar{J}(t) \right] \end{split}$$

 $\hookrightarrow$  non-diagonal kernels crucial for doubly-virtual case

● Another doubly-virtual complication: anomalous thresholds in time-like region Colangelo, MH, Procura, Stoffer arXiv:1309.6877

M. Hoferichter (Institute for Nuclear Theory)

Towards a data-driven analysis of HLbL scattering

# Subtraction functions

#### Omnès representation for S-wave

$$\begin{split} h_{0,++}(s) &= \Delta_{0,++}(s) + \Omega_0(s) \left[ \frac{1}{2} (s - s_+) a_+ (q_1^2, q_2^2) + \frac{1}{2} (s - s_-) a_- (q_1^2, q_2^2) + q_1^2 q_2^2 b(q_1^2, q_2^2) \right. \\ &+ \frac{s(s - s_+)}{2\pi} \int_{4M_\pi^2}^{\infty} ds' \frac{\sin \delta_0(s') \Delta_{0,++}(s')}{s'(s' - s_+)(s' - s) |\Omega_0(s')|} + \frac{s(s - s_-)}{2\pi} \int_{4M_\pi^2}^{\infty} ds' \frac{\sin \delta_0(s') \Delta_{0,++}(s')}{s'(s' - s_-)(s' - s) |\Omega_0(s')|} \\ &+ \frac{2q_1^2 q_2^2 s}{\pi} \int_{4M_\pi^2}^{\infty} ds' \frac{\sin \delta_0(s') \Delta_{0,00}(s')}{s'(s' - s_+)(s' - s_-) |\Omega_0(s')|} \right] \qquad s_{\pm} = q_1^2 + q_2^2 \pm 2\sqrt{q_1^2 q_2^2} \end{split}$$

• Inhomogeneities  $\Delta_{0,++}(s), \Delta_{0,00}(s)$  include left-hand cut

#### Subtraction functions

•  $b(q_1^2, q_2^2)$  and  $a_+(q_1^2, q_2^2) - a_-(q_1^2, q_2^2)$  multiply  $q_1^2 q_2^2$  and  $\sqrt{q_1^2 q_2^2}$ 

 $\hookrightarrow$  inherently doubly-virtual observables  $\Rightarrow$  need ChPT (or lattice)

- However:  $a(q_1^2, q_2^2) = (a_+(q_1^2, q_2^2) + a_-(q_1^2, q_2^2))/2$  fixed by singly-virtual measurements
  - $\hookrightarrow$  compare with chiral prediction, uncertainty estimates for the other functions are  $\neg \land \land \land$

• 1-loop result for arbitrary  $q_i^2$ , e.g.

$$\begin{aligned} \mathbf{a}^{\pi^{0}}(q_{1}^{2}, q_{2}^{2}) &= -\frac{M_{\pi}^{2}}{8\pi^{2}F_{\pi}^{2}(q_{1}^{2} - q_{2}^{2})^{2}} \left\{ q_{1}^{2} + q_{2}^{2} + 2\left(M_{\pi}^{2}(q_{1}^{2} + q_{2}^{2}) + q_{1}^{2}q_{2}^{2}\right)C_{0}(q_{1}^{2}, q_{2}^{2}) \right. \\ &+ q_{1}^{2}\left(1 + \frac{6q_{2}^{2}}{q_{1}^{2} - q_{2}^{2}}\right)\bar{J}(q_{1}^{2}) + q_{2}^{2}\left(1 - \frac{6q_{1}^{2}}{q_{1}^{2} - q_{2}^{2}}\right)\bar{J}(q_{2}^{2}) \right\} \end{aligned}$$

• Special case:  $q_1^2 = q_2^2 = 0$ 

$$a^{\pi^{\pm}}(0,0) = \frac{\overline{l_6} - \overline{l_5}}{48\pi^2 F_{\pi}^2} + \dots = \frac{M_{\pi}}{2\alpha} (\alpha_1 - \beta_1)^{\pi^{\pm}} \qquad b^{\pi^{\pm}}(0,0) = 0$$
$$a^{\pi^0}(0,0) = -\frac{1}{96\pi^2 F_{\pi}^2} + \dots = \frac{M_{\pi}}{2\alpha} (\alpha_1 - \beta_1)^{\pi^0} \qquad b^{\pi^0}(0,0) = -\frac{1}{1440\pi^2 F_{\pi}^2 M_{\pi}^2} + \dots$$

 $\hookrightarrow$  resum higher chiral orders into pion polarizabilities

## Subtraction functions: dispersive representation



Singly-virtual case: phenomenological representation with chiral constraints

- $\hookrightarrow$  parameters fixed from  $e^+e^- o \pi^0\pi^0\gamma$  (CMD2 and SND) Moussallam 2013
- **Dispersive representation**: imaginary part from  $2\pi$ ,  $3\pi$ , ...

 $\hookrightarrow$  analytic continuation from time-like to space-like kinematics

• Example:  $I = 2 \Rightarrow$  isovector photons  $\Rightarrow 2\pi \sim \rho$ 

$$\begin{aligned} a^{2}(q_{1}^{2},q_{2}^{2}) &= \alpha_{0} \Big[ \alpha^{2} + \alpha \Big( q_{1}^{2} \mathcal{F}^{\rho}(q_{1}^{2}) + q_{2}^{2} \mathcal{F}^{\rho}(q_{2}^{2}) \Big) + q_{1}^{2} q_{2}^{2} \mathcal{F}^{\rho}(q_{1}^{2}) \mathcal{F}^{\rho}(q_{2}^{2}) \\ \mathcal{F}^{\rho}(q^{2}) &= \frac{1}{\pi} \int_{4M_{\pi}^{2}}^{\infty} \mathrm{d}s \frac{q_{\pi\pi}^{3}(s) (F_{\pi}^{V}(s))^{*} \Omega_{1}(s)}{s^{3/2}(s - q^{2})} \qquad q_{\pi\pi}(s) = \sqrt{\frac{s}{4} - M_{\pi}^{2}} \end{aligned}$$

 $\hookrightarrow \alpha_0$  and  $\alpha$  can be determined from  $a^2(q^2, 0)$  alone!

Moussallam 2013

### Wick rotation: anomalous thresholds

Trajectory of the triangle anomalous thresholds for  $0 < q_1^2 < 4m^2$ 



M. Hoferichter (Institute for Nuclear Theory)