

# Naturalness, Supersymmetry and Light Higgsinos

XERXES TATA

University of Hawaii

H. Baer, V. Barger, P. Huang, D. Mickelson, A. Mustafayev, W. Sreethawong

- ★ SUSY solves the big hierarchy problem. Low scale physics does not have quadratic sensitivity to high scales if the low scale theory is embedded into a bigger framework with a high mass scale,  $\Lambda$ .
- ★ All talk about naturalness of weak scale SUSY models and associated fine-tuning has, at most, to do with logarithmic sensitivity to  $\Lambda$ .

Much discussion has revolved around the well-known (loop-corrected) minimization condition (written in terms of the parameters of the weak-scale theory),

$$\frac{M_Z^2}{2} = \frac{m_{H_d}^2 + \Sigma_d^d - (m_{H_u}^2 + \Sigma_u^u) \tan^2 \beta}{\tan^2 \beta - 1} - \mu^2,$$

requiring no large cancellations on the RHS.<sup>a</sup>

$$\Delta_{\text{EW}} = \max \left( \frac{m_{H_u}^2}{\frac{1}{2} M_Z^2} \frac{\tan^2 \beta}{\tan^2 \beta - 1}, \frac{\Sigma_u^u}{\frac{1}{2} M_Z^2} \frac{\tan^2 \beta}{\tan^2 \beta - 1}, \dots \right).$$

---

<sup>a</sup>In NUHM2, the choice of  $A_0$  that makes  $\Sigma_u^u$  small, simultaneously raises the Higgs boson mass.

$\Delta_{\text{EW}}$  knows nothing about the high scale physics, or the logs that we mentioned above. To see these, write

$$m_{H_u, H_d}^2 = m_{H_u, H_d}^2(\Lambda) + \delta m_{H_u, H_d}^2, \text{ etc.}$$

The logs are sitting in the  $\delta m_{\bullet}^2$  terms.

Define  $\Delta_{\text{HS}}$  analogously to  $\Delta_{\text{EW}}$ .

$\Delta_{\text{HS}}$  is a sensible measure of fine-tuning in that it knows about the high scale origins via the logs. If there are large cancellations between  $m_{\bullet}^2(\Lambda)$  and  $\delta m_{\bullet}^2(\Lambda)$ , the theory is regarded as fine-tuned. A VERY HIGH STANDARD!

However,  $\Delta_{\text{HS}}$  knows nothing about the correlations between various parameters that are present when the weak scale theory is derived from a high scale theory with fewer parameters.

To see these correlations, rewrite everything on the RHS in terms of the parameters of the theory defined at the scale  $\Lambda$ , *e.g.* (at one-loop)

$$\begin{aligned}
 -2\mu^2(m_{\text{weak}}) &= -2.18\mu^2 \\
 -2m_{H_u}^2(m_{\text{weak}}) &= 3.84M_3^2 + 0.32M_3M_2 + 0.047M_1M_3 - 0.42M_2^2 \\
 &\quad + 0.011M_2M_1 - 0.012M_1^2 - 0.65M_3A_t - 0.15M_2A_t \\
 &\quad - 0.025M_1A_t + 0.22A_t^2 + 0.004m_3A_b \\
 &\quad - 1.27m_{H_u}^2 - 0.053m_{H_d}^2 \\
 &\quad + 0.73m_{Q_3}^2 + 0.57m_{U_3}^2 + 0.049m_{D_3}^2 - 0.052m_{L_3}^2 + 0.053m_{E_3}^2 \\
 &\quad + 0.051m_{Q_2}^2 - 0.11m_{U_2}^2 + 0.051m_{D_2}^2 - 0.052m_{L_2}^2 + 0.053m_{E_2}^2 \\
 &\quad + 0.051m_{Q_1}^2 - 0.11m_{U_1}^2 + 0.051m_{D_1}^2 - 0.052m_{L_1}^2 + 0.053m_{E_1}^2,
 \end{aligned}$$

In a theory with universal scalar masses  $m_0$  and universal gaugino masses  $m_{1/2}$  and a universal  $A$ -parameter  $A_0$  this collapses to,

$$-2m_{H_u}^2(m_{\text{weak}}) = 3.78m_{1/2}^2 - 0.82A_0m_{1/2} + 0.22A_0^2 + 0.013m_0^2$$

.

We can substitute this along with analogous expression for  $m_{H_d}^2$  (which usually makes a small effect) into the  $M_Z^2$  expression that we had, and again require no large cancellations.

Clearly, the inclusion of the correlations make an important difference as they allow some cancellations between  $m_{\bullet}^2(\Lambda)$  and the  $\delta m_{\bullet}^2$  without counting to the fine-tuning.

Related to the often-used Barbieri-Guidice  $\Delta_{BG}$  measure first introduced by EENZ.

$$\Delta_{HS} > \Delta_{BG} > \Delta_{EW}.$$

## What use is $\Delta_{EW}$ if it does not know about logs?

Imagine a high scale theory in which the combination  $m_{H_u}^2(\Lambda) + \delta m_{H_u}^2$  is automatically small. In such a theory,  $\Delta_{EW}$  is a perfectly sensible fine-tuning measure!

- ★ In the NUHM2 model – mSUGRA +  $m_{H_u}^2, m_{H_d}^2$  as independent parameters – this is guaranteed for special values of  $m_{H_u}^2$ , so “all we have to do” is find the theory that leads to this value of  $m_{H_u}^2$ !
- ★ The FP/HB region of mSUGRA and its generalizations if  $\Lambda \sim M_P$  has partial automatic cancellations.
- ★ Mixed-modulus-anomaly-mediated-SUSY-breaking models aka mirage-mediation models, for special values of  $\alpha$ .

## Properties of $\Delta_{EW}$ .

- ★  $\Delta_{EW}$  is essentially determined by the SUSY spectrum.
- ★ If  $\Delta_{EW}$  is large, the underlying theory that leads to the spectrum will be fine-tuned. A small  $\Delta_{EW}$  does not imply the theory is not fine-tuned, but leaves open the possibility of finding such a theory.

Low  $\Delta_{EW} \implies$  low  $|\mu|$ , but squarks (including stops) may be much heavier.

Many aspects of the phenomenology depend just on the spectrum, so this can be investigated even without knowledge of the underlying high scale theory that leads to low fine-tuning. Beware though of pheno implications that depend on correlations in the spectrum.

We think low  $|\mu|$  more basic to fine-tuning considerations than light stops. This feature is hidden by many analyses of fine-tuning.

Very generally, light higgsinos are a necessary feature of models with low fine-tuning.



## Phenomenology with light higgsinos.

- ★ Higgsino-like states  $\widetilde{W}_1^\pm$ ,  $\widetilde{Z}_2$ ,  $\widetilde{Z}_1$  must be present with masses  $\sim |\mu|$ , and generically small splittings.
- ★ If  $|M_{1,2}|$  also happen to be comparable to  $|\mu|$ , these states would be easy to access at the LHC via  $\widetilde{W}_1 \widetilde{Z}_2$  production, or at a \*LC via  $\widetilde{W}_1 \widetilde{W}_1$ ,  $\widetilde{Z}_1 \widetilde{Z}_2$  and  $\widetilde{Z}_2 \widetilde{Z}_2$  production. Heavier -inos may also be accessible.
- ★ In the generic case, the small mass gap may make it difficult to see direct higgsino production signals at the LHC because decay products are very soft.
- ★ Need careful investigation of how small a mass gap we can probe to discriminate signal from two-photon production of heavy flavours at the \*LC. Early studies of the HB/FP region of mSUGRA indicative of good news using specialized cuts.
- ★ A novel signal is also possible at the LHC if  $|M_2| \lesssim 0.8 - 1$  TeV, something that is possible but not compulsory for low  $\Delta_{EW}$ .



## Light higgsinos at the LHC

Unified gaugino masses for definiteness,  $\mu = 150$  GeV




$\approx 10$  fb  $\tilde{W}_2\tilde{Z}_4$  cross section out to  $m_{1/2} \sim 1$  TeV

Decays of the parent  $\tilde{W}_2$  and  $\tilde{Z}_4$  that lead to  $W$  boson pairs give the same sign 50% of the time.

Require exactly two isolated, same sign dileptons with  $|p_T(\ell)| > 20$  GeV, and no tagged  $b$ -jets (60% eff.)  $m_T^{\min} > 125$  GeV cut removes  $WZ$  and  $t\bar{t}$  backgrounds.

NUHM2:  $m_0=5$  TeV,  $A_0=-1.6m_0$ ,  $\tan\beta=15$ ,  $\mu=150$  GeV,  $m_A=1$  TeV



arXiv:1303.3816

Reach to  $m_{1/2} \sim 680$  (1000) GeV for 100 (1000)  $\text{fb}^{-1}$ . In low  $\mu$  models, this is better than the canonical trilepton reach of 400 (500) GeV for 300 (1000)  $\text{fb}^{-1}$ .

In models with gaugino mass unification, this channel offers a better reach than the usual  $\tilde{g}\tilde{g}$  channel. More importantly, this is an independent search channel.

## Final remarks

- ★ Obituaries of SUSY are premature. Models with modest electroweak fine-tuning ( $\Delta_{EW} = 10 - 30$ ) and correct Higgs boson mass exist. These satisfy our necessary criteria for fine-tuning. **Eagerly awaiting LHC14.**
- ★ Small  $|\mu|$  is a fundamental and necessary (but not sufficient) criterion for low fine-tuning.
- ★  $\Delta_{EW}$  is “directly” measurable (in principle) so we can tell for sure that a given spectrum is fine-tuned if  $\Delta_{EW}$  turns out to be large.
- ★ Light higgsino scenarios cannot saturate the total CDM; nonetheless there is enough higgsino DM fraction that will reveal itself in direct and indirect DM searches. (arXiv:1303.3816, Baer, Barger, Mickelson)
- ★ Novel **SS dilepton signal with low hadronic activity possible at the LHC.**
- ★ A \*LC with  $\sqrt{s} \gtrsim (500)700$  GeV could be a discovery machine for light higgsinos for  $\Delta_{EW} \lesssim (15)30$ .