

Microstructure and Particle-Phase Stress in a Dense Suspension

Jim Jenkins, Cornell University


Luigi La Ragione, Politecnico di Bari

Predict microstructure and particle stresses in
slow, steady, pure straining of a dense, single layer
of spheres in a viscous fluid

Singh & Nott, J Fluid Mech. **412**, 279 (2000)

Jenkins & La Ragione, J. Fluid Mech. **763**, 218 (2015)

Pure Straining



Area fraction: ν

Average number of near contacts: k

Short-range repulsive force: F_0

Kinematics

Treat BA exactly

Relative velocity of **centers** of pair BA

$$v_{\alpha}^{(BA)} = \frac{d}{dt} s^{(BA)} \hat{d}_{\alpha}^{(BA)} + 2a \frac{d}{dt} \theta^{(BA)} \hat{t}_{\alpha}^{(BA)}$$

Relative velocity of **points of near contact** of BA

$$v_{\alpha}^{(BA)} + a \left(\omega^{(A)} + \omega^{(B)} \right) \hat{t}_{\alpha}^{(BA)}$$

Treat nA on average

Relative velocity of **centers** of pair nA

$$v_i^{(nA)} = 2a D_{\alpha\beta} \hat{d}_{\beta}^{(nA)}$$

Relative velocity of **points of near contact** of nA

$$v_{\alpha}^{(nA)} + a \omega^{(A)} \hat{t}_{\alpha}^{(nA)}$$

Forces of interaction

Jeffrey & Onishi (1984), Jeffrey (1992)

Pair BA

$$F_{\alpha}^{(BA)} = 6\pi\mu a K_{\alpha\beta}^{(BA)} v_{\beta}^{(BA)} - \frac{F_0}{s^{(BA)}} \hat{d}_{\alpha}^{(BA)} - 9.54\pi\mu a^2 \left(\hat{t}_{\beta}^{(BA)} D_{\beta\gamma} \hat{d}_{\gamma}^{(BA)} \right) \hat{t}_{\alpha}^{(BA)} \\ + \pi\mu a^2 \left[\ln\left(\frac{a}{s^{(BA)}}\right) - 0.96 \right] \omega^{(A)} \hat{t}_{\alpha}^{(BA)} + \pi\mu a^2 \ln\left(\frac{a}{s^{(BA)}}\right) \omega^{(B)} \hat{t}_{\alpha}^{(BA)}$$

$$K_{\alpha\beta}^{(BA)} = \frac{1}{4} \frac{a}{s^{(BA)}} \hat{d}_{\alpha}^{(BA)} \hat{d}_{\beta}^{(BA)} + \frac{1}{6} \ln\left(\frac{a}{s^{(BA)}}\right) \hat{t}_{\alpha}^{(BA)} \hat{t}_{\beta}^{(BA)}$$

Pairs nA

$$F_{\alpha}^{(nA)} = 12\pi\mu a^2 K_{\alpha\beta}^{(nA)} D_{\beta\gamma} \hat{d}_{\gamma}^{(nA)} + \pi\mu a^2 \left[\ln\left(\frac{a}{\bar{s}}\right) - 0.96 \right] \omega^{(A)} \hat{t}_{\alpha}^{(nA)} \\ - \frac{F_0}{\bar{s}} \hat{d}_{\alpha}^{(nA)} + 2\pi\mu a^2 \left[\ln\left(\frac{1}{\bar{s}}\right) - 0.96 \right] \left(\hat{t}_{\beta}^{(nA)} D_{\beta\gamma} \hat{d}_{\gamma}^{(nA)} \right) \hat{t}_{\alpha}^{(nA)}$$

$$K_{\alpha\beta}^{(nA)} = \frac{1}{4} \frac{a}{\bar{s}} \hat{d}_{\alpha}^{(nA)} \hat{d}_{\beta}^{(nA)} + \frac{1}{6} \ln\left(\frac{a}{\bar{s}}\right) \hat{t}_{\alpha}^{(nA)} \hat{t}_{\beta}^{(nA)}$$

Force Balance, Particle A

$$F_{\alpha}^{(BA)} + \sum_{n \neq B}^{N^{(A)}} F_{\alpha}^{(nA)} = 0$$

$$F_{\alpha}^{(AB)} = -F_{\alpha}^{(BA)}, \quad \hat{d}_{\alpha}^{(AB)} = -\hat{d}_{\alpha}^{(BA)}, \quad \hat{t}_{\alpha}^{(AB)} = -\hat{t}_{\alpha}^{(BA)}$$

Difference of force balances

Normal component

$$0 = 3\pi\mu a \frac{a}{s^{(BA)}} \frac{d}{dt} s^{(BA)} - 2 \frac{F_0}{s^{(BA)}} \\ + 6\pi\mu a^2 \frac{a}{\bar{S}} \hat{d}_{\alpha}^{(BA)} J_{\alpha\beta\gamma} D_{\beta\gamma} - 2 \frac{F_0}{\bar{S}} Y_{\alpha} \hat{d}_{\alpha}^{(BA)}$$

Tangential component

$$0 = 4\pi\mu a^2 \left[\ln\left(\frac{a}{s^{(BA)}}\right) + 3.84 \right] \frac{d}{dt} \theta^{(BA)} - 19\pi\mu a^2 \hat{t}_{\alpha}^{(BA)} D_{\alpha\beta} \hat{d}_{\beta}^{(BA)} \\ + \pi\mu a^2 \left[2 \ln\left(\frac{a}{s^{(BA)}}\right) - 0.96 \right] \bar{S} + 6\pi\mu a^2 \frac{a}{\bar{S}} \hat{t}_{\alpha}^{(BA)} J_{\alpha\beta\gamma} D_{\beta\gamma} \\ (\bar{S} = \omega^A + \omega^B)$$

Moment Balance, Particle A

$$\epsilon_{\alpha\beta} F_{\alpha}^{(BA)} \hat{d}_{\beta}^{(BA)} + \epsilon_{\alpha\beta} \sum_{n \neq B}^{N(A)} F_{\alpha}^{(nA)} \hat{d}_{\beta}^{(nA)} = 0$$

$$(\epsilon_{12} = -\epsilon_{21} = 1, \epsilon_{11} = \epsilon_{22} = 0)$$

Sum of Moment Balances

$$0 = 2 \left[\ln \left(\frac{a}{s^{(BA)}} \right) + 3.84 \right] \frac{d}{dt} \theta^{(BA)} - 9.54 \hat{t}_{\alpha}^{(BA)} D_{\alpha\beta} \hat{d}_{\beta}^{(BA)}$$

$$+ \left[\ln \left(\frac{a}{s^{(BA)}} \right) + \frac{k-1}{2} \ln \left(\frac{a}{\bar{s}} \right) \right] S + 2 \left[\ln \left(\frac{a}{\bar{s}} \right) - 0.96 \right] \epsilon_{\beta\gamma} A_{\alpha\beta} D_{\alpha\gamma}$$

Structural Sums

Isotropic representations

$$Y_{\alpha} \equiv \overline{\sum_{n \neq B}^{N(A)} \hat{d}_{\alpha}^{(nA)}} = \xi \hat{d}_{\alpha}^{(BA)}$$

$$A_{\alpha\beta} \equiv \overline{\sum_{n \neq B}^{N(A)} \hat{d}_{\alpha}^{(nA)} \hat{d}_{\beta}^{(nA)}} = \beta_1 \delta_{\alpha\beta} + \beta_2 \hat{d}_{\alpha}^{(BA)} \hat{d}_{\beta}^{(BA)}$$

$$\begin{aligned} J_{\alpha\beta\gamma} &\equiv \overline{\sum_{n \neq B}^{N(A)} \hat{d}_{\alpha}^{(nA)} \hat{d}_{\beta}^{(nA)} \hat{d}_{\gamma}^{(nA)}} \\ &= \left[\alpha_1 \hat{d}_{\alpha}^{(BA)} \hat{d}_{\beta}^{(BA)} \hat{d}_{\gamma}^{(BA)} \right. \\ &\quad \left. + \alpha_2 \left(\delta_{\alpha\beta} \hat{d}_{\gamma}^{(BA)} + \delta_{\alpha\gamma} \hat{d}_{\beta}^{(BA)} + \delta_{\beta\gamma} \hat{d}_{\alpha}^{(BA)} \right) \right] \end{aligned}$$

Averages

$$\frac{\bar{s}}{a} = \frac{\sqrt{\pi}}{8\mathbf{v}g_0} \quad g_0 = \frac{16 - 7\mathbf{v}}{16(1 - \mathbf{v})^2}$$

Isotropic distribution function

$$\overline{\Psi_{\alpha.. \gamma}} = 2 \int_0^{\pi} I(\theta') \hat{d}_{\alpha} \dots \hat{d}_{\gamma} d\theta'$$

$$I(\theta') = \begin{cases} 0, & 0 < \theta' < \pi/3 \\ 3(\mathbf{k} - 1) / 4\pi, & \pi/3 < \theta' < \pi \end{cases}$$

$$\alpha_1 = 0, \quad \alpha_2 = -\frac{3\sqrt{3}(\mathbf{k} - 1)}{16\pi} \equiv \alpha$$

$$\beta_1 = \frac{(\mathbf{k} - 1)}{2} + \frac{3\sqrt{3}(\mathbf{k} - 1)}{16\pi}, \quad \beta_2 = -\frac{3\sqrt{3}(\mathbf{k} - 1)}{8\pi} = 2\alpha$$

$$\xi = -\frac{3\sqrt{3}(\mathbf{k} - 1)}{4\pi} = 4\alpha$$

Pure Straining

$$\mathbf{D}_{\alpha\beta} = \begin{bmatrix} \dot{\gamma} & 0 \\ 0 & -\dot{\gamma} \end{bmatrix} \quad \hat{\mathbf{d}}_{\alpha}^{(\text{BA})} = (\sin \theta, \cos \theta)$$

$$\hat{\mathbf{d}}_{\alpha}^{(\text{BA})} \mathbf{D}_{\alpha\beta} \hat{\mathbf{d}}_{\beta}^{(\text{BA})} = -\dot{\gamma} \cos 2\theta, \quad \hat{\mathbf{t}}_{\alpha}^{(\text{BA})} \mathbf{D}_{\alpha\beta} \hat{\mathbf{d}}_{\beta}^{(\text{BA})} = \dot{\gamma} \sin 2\theta$$

Lengths dimensionless by a and $F \equiv F_0 / (\pi \mu a^3 \dot{\gamma})$

$$\frac{1}{s} \frac{ds}{d\gamma} = \frac{2}{3} F \left(\frac{1}{s} + 4 \frac{\alpha}{\bar{s}} \right) + 4 \frac{\alpha}{\bar{s}} \cos 2\theta$$

$$\left[\ln \left(\frac{1}{s} \right) + 3.84 \right] \frac{d\theta}{d\gamma} = \left(4.77 - 3\alpha \frac{1}{\bar{s}} \right) \sin 2\theta - \frac{1}{2} \ln \left(\frac{1}{s} \right) s$$

Tangential force difference - Sum moments

$$s = - \frac{12\alpha}{(4\alpha - k + 1)} \frac{1/\bar{s}}{\ln(1/\bar{s}) - 0.96} \sin 2\theta$$

Contact Distribution

$A(\theta)d\theta$: number of contacts within $d\theta$

$$A(\theta) \frac{d\theta}{d\gamma} = \text{constant} \quad \text{and} \quad \int_0^{\pi/2} A(\theta) d\theta = \frac{k}{4}$$

Knowledge of k determines the constant of integration.

Implement as differential equations

$$I(\theta) \equiv \int_0^\theta A(\theta') d\theta'$$

$$\frac{dI}{d\theta} = A, \quad I(0) = 0, \quad I(\pi/2) = k/4$$

$$\frac{dA}{d\theta} = -\frac{A}{\dot{\theta}} \frac{d\dot{\theta}}{d\theta} \quad \frac{d\dot{\theta}}{d\theta} = \frac{\partial \dot{\theta}}{\partial \theta} + \frac{\partial \dot{\theta}}{\partial s} \frac{ds}{d\theta}$$

Differential Equations

$$\frac{ds}{d\theta} = \frac{ds/d\gamma}{d\theta/d\gamma} \equiv f[s, \theta; \bar{s}(v), k]$$

$$\frac{dA}{d\theta} = \frac{A}{d\theta/d\gamma} \left[\frac{\partial}{\partial \theta} \left(\frac{d\theta}{d\gamma} \right) + \frac{\partial}{\partial s} \left(\frac{d\theta}{d\gamma} \right) \frac{ds}{d\theta} \right]$$

$$\frac{dI}{d\theta} = A$$

$$\frac{d\gamma}{d\theta} = \frac{1}{d\theta/d\gamma}$$

Boundary Conditions


$$0 \leq x \leq 1 \quad \theta = \theta_0 + (\theta_1 - \theta_0)x$$

$$s(\theta_0) = 0.003 \quad s(\theta_1) = 0.003$$

$$I(\theta_0) = 0 \quad I(\theta_1) = k/4$$


$$\gamma(\theta_0) = 0 \quad \gamma(\theta_1) = 1.1$$

Separation




The repulsive force creates asymmetry about $\pi/4$.

Inverse Separation



The asymmetry of the separation clarified.


Angular Velocity



$$A(\theta) \frac{d\theta}{d\gamma} = \text{constant}$$


Circumferential Distribution

$$k = 3.0, \gamma = 1.1, s_0 = s_1 = 0.003$$



The repulsive force shifts the anisotropy due to the flow, symmetric about $\pi/4$, to a distribution that is asymmetric.

Circumferential Distribution



Asymmetry about $\pi/4$ in both the distribution and interval.

Stress (dimensional)

$$T_{\alpha\beta} = na \int_0^{2\pi} A(\theta) F_{\alpha}^{(BA)} \hat{d}_{\beta}^{(BA)} d\theta$$

n: number/area

$$F_{\alpha}^{(BA)} = - \sum_{n \neq B}^{N^{(A)}} F_{\alpha}^{(nA)}$$

$$t_{\alpha\beta} = \frac{T_{\alpha\beta}}{a\mu\dot{\gamma}}, \quad n = \frac{v}{\pi a^2}, \quad F = \frac{F_0}{\pi a^3 \mu \dot{\gamma}}$$

$$\begin{aligned} t_{\alpha\beta} = & v \frac{\alpha}{\bar{s}} \int_0^{2\pi} A(\theta) (4F + 6 \cos 2\theta) \hat{d}_{\alpha} \hat{d}_{\beta} d\theta \\ & + v \left(9.54 - \frac{6}{\bar{s}} \right) \alpha \int_0^{2\pi} A(\theta) \sin 2\theta \hat{t}_{\alpha} \hat{d}_{\beta} d\theta \\ & + v \frac{12\alpha}{(3\alpha - k + 1) \ln(1/\bar{s})} \frac{1/\bar{s}}{\ln(1/\bar{s})} \int_0^{2\pi} A(\theta) \ln\left(\frac{1}{s}\right) \sin 2\theta \hat{t}_{\alpha} \hat{d}_{\beta} d\theta \end{aligned}$$

$$\hat{d}_{\alpha}^{(AB)} = (\sin \theta, \cos \theta), \quad \hat{t}_{\alpha}^{(AB)} = (\cos \theta, -\sin \theta)$$

Pressure

$$p \equiv -\frac{1}{2}(t_{xx} + t_{yy})$$

$$p = -4\nu \frac{\alpha}{\bar{s}} \int_0^{\pi/2} A(\theta) (2F + 3\cos 2\theta) d\theta$$

Shear stress


$$\tau \equiv \frac{1}{2}(t_{xx} - t_{yy})$$

$$\tau = -4\nu \frac{\alpha}{\bar{s}} \int_0^{\pi/2} A(\theta) (2F + 3\cos 2\theta) \cos 2\theta d\theta$$

$$+ \nu \left(4.77 - \frac{3\alpha}{\bar{s}} \right) \int_0^{2\pi} A(\theta) \sin^2 2\theta d\theta$$


$$+ \nu \frac{6\alpha}{(3\alpha - k + 1)} \frac{1/\bar{s}}{\ln(1/\bar{s})} \int_0^{2\pi} A(\theta) \ln\left(\frac{1}{s}\right) \sin^2 2\theta d\theta$$

Stresses



Asymmetry about $\pi/4$ creates particle pressure.

Initial/Final Separation



The initial/final separation of the “typical” trajectory is where the shear stress exhibits a maximum (0.003).

Conclusions

A rough implementation of **particle equilibrium** for particles assumed to **interact as pairs through points of near contact** produces expressions for the **normal and tangential components of their relative velocities as functions of their orientation with respect to the flow.**

From these velocities, the separation of neighboring particles and their **circumferential distribution can be determined as functions of orientation.**

The existence of **asymmetries in the circumferential distribution of particles** influences the shear stress and **produces a pressure** associated with viscous interactions between particle pairs at points of near contact.

Include elasticity and friction

Asperity height $s_0/2$, Stiffness: k_0 , Sliding friction: f

If $s > s_0$

$$F_{\alpha}^{(AB)} \doteq 6\pi\mu a K_{\alpha\beta}^{(AB)} v_{\beta}^{(AB)} - \frac{F_0}{s^{(AB)}} \hat{d}_{\alpha}^{(AB)} - 9.54\pi\mu a^2 \left(\hat{t}_{\beta}^{(AB)} D_{\beta\gamma} \hat{d}_{\gamma}^{(AB)} \right) \hat{t}_{\alpha}^{(AB)}$$

else

$$F_{\alpha}^{(AB)} \doteq 6\pi\mu a K_{\alpha\beta}^{(AB)} v_{\beta}^{(AB)} - \frac{F_0}{s^{(AB)}} \hat{d}_{\alpha}^{(AB)} - 9.54\pi\mu a^2 \left(\hat{t}_{\beta}^{(AB)} D_{\beta\gamma} \hat{d}_{\gamma}^{(AB)} \right) \hat{t}_{\alpha}^{(AB)} \\ - k_0(s_0 - s) \hat{d}_{\alpha}^{(AB)} - f k_0(s_0 - s) \hat{t}_{\alpha}^{(AB)}$$

Coefficient of sliding friction: f

$$K_{\alpha\beta}^{(BA)} = \frac{1}{4} \frac{a}{s^{(BA)}} \hat{d}_{\alpha}^{(BA)} \hat{d}_{\beta}^{(BA)} + \frac{1}{6} \ln \left(\frac{a}{s^{(BA)}} \right) \hat{t}_{\alpha}^{(BA)} \hat{t}_{\beta}^{(BA)}$$

$$F_{\alpha}^{(nA)} \doteq \frac{3}{\bar{s}} \pi\mu a^2 \left(\hat{d}_{\beta}^{(nA)} D_{\beta\gamma} \hat{d}_{\gamma}^{(nA)} \right) \hat{d}_{\alpha}^{(AB)} \\ - \frac{F_0}{\bar{s}} \hat{d}_{\alpha}^{(nA)} + \pi\mu a^2 \left[2 \ln \left(\frac{1}{\bar{s}} \right) - 1.92 \right] \left(\hat{t}_{\beta}^{(AB)} D_{\beta\gamma} \hat{d}_{\gamma}^{(AB)} \right) \hat{t}_{\alpha}^{(AB)}$$

Normal:

$$0 = \frac{1}{s^{(BA)}} \frac{d}{dt} s^{(BA)} + \frac{4\alpha}{\bar{s}} \hat{d}_{\beta}^{(BA)} D_{\beta\gamma} \hat{d}_{\gamma}^{(BA)} - \frac{2F_0}{3\pi\mu a^2} \left(\frac{1}{s^{(BA)}} + \frac{4\alpha}{\bar{s}} \right) - \frac{2k_0}{3\pi\mu a^2} \left[\Theta(s_0 - s^{(BA)})(s_0 - s^{(BA)}) + 4\alpha\Theta(s_0 - \bar{s})(s_0 - \bar{s}) \right]$$

Tangential:

$$0 = 12 \left[\frac{1}{6} \ln \left(\frac{1}{s^{(BA)}} \right) + 0.64 \right] \frac{d}{dt} \theta^{(BA)} - 9.54 \hat{t}_{\beta}^{(BA)} D_{\beta\gamma} \hat{d}_{\gamma}^{(BA)} + 2\alpha \left[2 \ln \left(\frac{1}{\bar{s}} \right) - 1.92 \right] \hat{t}_{\beta}^{(BA)} D_{\beta\gamma} \hat{d}_{\gamma}^{(BA)} + \frac{6\alpha}{\bar{s}} \hat{t}_{\beta}^{(BA)} D_{\beta\gamma} \hat{d}_{\gamma}^{(BA)} - \frac{k_0 f}{\pi\mu a^2} \left[\Theta(s_0 - s^{(BA)})(s_0 - s^{(BA)}) + 2\alpha\Theta(s_0 - \bar{s})(s_0 - \bar{s}) \right]$$

Dimensionless:

$$0 = \frac{1}{s} \frac{ds}{dt} - \frac{2\alpha}{\bar{s}} \cos 2\theta - \frac{2F}{3} \left(\frac{1}{s} + \frac{4\alpha}{\bar{s}} \right) - \frac{2k}{3} \Theta(s_0 - s)(s_0 - s) - \left[\ln \left(\frac{1}{s} \right) + 3.84 \right] \frac{d\theta}{dt} + \alpha \left[\ln \left(\frac{1}{\bar{s}} \right) - 0.96 \right] \sin 2\theta + \left(\frac{3\alpha}{2\bar{s}} - 4.77 \right) \sin 2\theta - \frac{kf}{4} \Theta(s_0 - s)(s_0 - s)$$

Brady

If $s > s_0$

$$F_{\alpha}^{(AB)} \doteq 6\pi\mu a K_{\alpha\beta}^{(AB)} v_{\beta}^{(AB)} - \frac{F_0}{s^{(AB)}} \hat{d}_{\alpha}^{(AB)} \\ - 9.54\pi\mu a^2 \left(\hat{t}_{\beta}^{(AB)} D_{\beta\gamma} \hat{d}_{\gamma}^{(AB)} \right) \hat{t}_{\alpha}^{(AB)} - \pi\mu a \ln \left(\frac{a}{s^{(BA)}} \right) \hat{t}_{\alpha}^{(BA)}$$

else

$$F_{\alpha}^{(AB)} \doteq 6\pi\mu a K_{\alpha\beta}^{(AB)} v_{\beta}^{(AB)} - \frac{F_0}{s^{(AB)}} \hat{d}_{\alpha}^{(AB)} \\ - 9.54\pi\mu a^2 \left(\hat{t}_{\beta}^{(AB)} D_{\beta\gamma} \hat{d}_{\gamma}^{(AB)} \right) \hat{t}_{\alpha}^{(AB)} - \frac{3}{2} \pi\mu a \frac{a}{s^{(BA)}} \hat{t}_{\alpha}^{(AB)}$$

$$T_{\alpha\beta} = na \int_0^{2\pi} A(\theta) F_{\alpha}^{(BA)} \hat{d}_{\beta}^{(BA)} d\theta$$

$$t_{\alpha\beta} = \frac{T_{\alpha\beta}}{a\mu\dot{\gamma}}, \quad n = \frac{v}{\pi a^2}, \quad F = \frac{F_0}{\pi a^3 \mu \dot{\gamma}}$$

$$\begin{aligned} t_{\alpha\beta} &= v \frac{\alpha}{\bar{s}} \int_0^{2\pi} A(\theta) (4F + 6 \cos 2\theta) \hat{d}_{\alpha} \hat{d}_{\beta} d\theta \\ &+ v \left(9.54 - \frac{6}{\bar{s}} \right) \int_0^{2\pi} A(\theta) \sin 2\theta \hat{t}_{\alpha} \hat{d}_{\beta} d\theta \\ &+ v \frac{12\alpha}{(3\alpha - k + 1) \ln(1/\bar{s})} \int_0^{2\pi} A(\theta) \ln\left(\frac{1}{s}\right) \sin 2\theta \hat{t}_{\alpha} \hat{d}_{\beta} d\theta \end{aligned}$$

$$\hat{d}_{\alpha}^{(BA)} D_{\alpha\beta} \hat{d}_{\beta}^{(BA)} = -\dot{\gamma} \cos 2\theta \quad \hat{t}_{\alpha}^{(BA)} D_{\alpha\beta} \hat{d}_{\beta}^{(BA)} = \dot{\gamma} \sin 2\theta$$

$$\begin{aligned} t_{\alpha\beta} &= v \frac{\alpha}{\bar{s}} \int_0^{2\pi} A(\theta) \left(4F - 6 \hat{d}_{\gamma}^{(BA)} \tilde{D}_{\gamma\mu} \hat{d}_{\mu}^{(BA)} \right) \hat{d}_{\alpha} \hat{d}_{\beta} d\theta \\ &+ v \left(9.54 - \frac{6}{\bar{s}} \right) \int_0^{2\pi} A(\theta) \hat{t}_{\gamma}^{(BA)} \tilde{D}_{\gamma\mu} \hat{d}_{\mu}^{(BA)} \hat{t}_{\alpha} \hat{d}_{\beta} d\theta \\ &+ v \frac{12\alpha}{(3\alpha - k + 1) \ln(1/\bar{s})} \int_0^{2\pi} A(\theta) \ln\left(\frac{1}{s}\right) \hat{t}_{\gamma}^{(BA)} \tilde{D}_{\gamma\mu} \hat{d}_{\mu}^{(BA)} \hat{t}_{\alpha} \hat{d}_{\beta} d\theta \end{aligned}$$

$$\tilde{D}_{\alpha\beta} \equiv \dot{\gamma}^{-1} D_{\alpha\beta}$$

$$\begin{aligned}
t_{\alpha\beta} = & \left[v \frac{\alpha}{\bar{s}} \int_0^{2\pi} A(\theta) \left(4F \hat{d}_\alpha \hat{d}_\beta - 6 \hat{d}_\alpha \hat{d}_\beta \hat{d}_\gamma^{(BA)} \hat{d}_\mu^{(BA)} \right) d\theta \right. \\
& + v \left(9.54 - \frac{6}{\bar{s}} \right) \int_0^{2\pi} A(\theta) \hat{t}_\gamma^{(BA)} \hat{d}_\mu^{(BA)} \hat{t}_\alpha \hat{d}_\beta d\theta \\
& \left. + v \frac{12\alpha}{(3\alpha - k + 1)} \frac{1/\bar{s}}{\ln(1/\bar{s})} \int_0^{2\pi} A(\theta) \ln\left(\frac{1}{s}\right) \hat{t}_\gamma^{(BA)} \hat{d}_\mu^{(BA)} \hat{t}_\alpha \hat{d}_\beta d\theta \right] \tilde{D}_{\gamma\mu}
\end{aligned}$$

$$\text{with } \hat{t}_\alpha = \varepsilon_{\alpha\beta} \hat{d}_\beta,$$